

OVERVIEW

The Riemann Hypothesis

The three Millennium problems: RH, NSE, YME

Dr. Klaus Braun
August 11, 2018

Albert Einstein, *"we can't solve problems by using the same kind of thinking we used when we created them"*,

Wolfgang E. Pauli, *"all things reach the one who knows how to wait"*.

This homepage addresses the following three Millennium problems (resp. links to corresponding homepages):

- A. The Riemann Hypothesis (RH)
- B. The 3D-Navier-Stokes equations (NSE)
- C. The Yang-Mills equations (YME)

A helicopter view

From a helicopter point of view there is a common denominator of all solution concepts of this homepage: it is about a common mathematical frame to govern the "infinitesimal small" with respect to truly infinitesimal small "elements" and related "functions" and truly geometrical (i.e. equipped with an inner product) "function spaces", enabling a truly geometrical mathematical modelling framework with corresponding operators (including well defined domains and ranges).

Regarding the RH the "infinitesimal small" is about the challenge to represent the entire Zeta function as a (Mellin-) transform of a self-adjoint operator ([EdH] 10.3). The Hilbert transform is the proposed tool to build a self-adjoint operator enabling the Berry-Keating (Hilbert-Polya) conjecture, because of its property that any Hilbert transformed function has a vanishing constant Fourier term (see also Polya's Bessel function based alternative entire Zeta function representation, [EdH], 12.5).

Regarding the NSE and YME, beside the n-dimensional counterpart of the Hilbert transform, the Riesz transforms, the two applied central "objects" are the well-established "differentials" and the distributional "Hilbert scale" concept enabling Pseudo-Differential and Fourier multiplier (weak and strong singular integral) equations (Calderón) and the related (Stieltjes integral like) spectral representation of Hermitian operators.

In the context of the newly proposed "energy-space" $H_{1/2} = H_1 + H_1^-$ we also refer to the Bose-Einstein Condensation (BEC), where below the critical temperature T_c BEC "normal gas" particles coexist in equilibrium with "condensed" particles. Unlike a liquid droplet in a gas, here the "condensed" particles are not separated in space ($H_{-1/2} = H_0 + H_0^-$) from normal particles. Instead they are separated in standard momentum space H_1 , which is a closed, compactly embedded subspace of the newly proposed "energy-space" $H_{1/2}$. The condensed standard particles all occupy a single quantum state of zero momentum, while normal standard particles all have finite momentum with respect to the H_1 - norm.

The challenge to represent to entire Zeta function as a (Mellin-) transform of a self-adjoint operator is explained in [EdH], 10.3:

Let $\gamma(s)$, $Z(s)$, $z(s)$ denote the Gamma function, the entire Zeta function ([EdH] 1.8 (1)) and the Zeta function ([EdH] 1.4 (3))

$$z(s) = \frac{\gamma(1-s)}{2\pi i} \int_{+\infty}^{+\infty} \frac{(-x)^s dx}{e^{x-1} x},$$

which is analytic at all points of the complex s-plane except for a simple pole at $s = 1$. It is equal to the Dirichlet function

$$z(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \text{ for } \operatorname{Re}(s) > 1 .$$

Let further denote $f(x) := e^{-\pi x^2}$ denote the Gaussian function (with mean value $f(0) = 1$) and

$$M[f](s) := \int_0^{\infty} f(x) x^s \frac{dx}{x} = \frac{1}{2} \gamma\left(\frac{s}{2}\right) \pi^{-s/2} , \quad G(x) := \sum_{n=1}^{\infty} f(nx).$$

Then it holds ([EdH] 1.6 (5), 12.5)

$$(*) \quad \frac{1}{2} \gamma\left(\frac{s}{2}\right) \pi^{-s/2} Z(s) = \frac{1}{2} \gamma\left(\frac{1-s}{2}\right) \pi^{-\frac{1-s}{2}} Z(1-s)$$

$$(**) \quad Z(s) = \frac{1}{2} \int_0^{\infty} x^{1-s} \frac{d}{dx} \left(x^2 \frac{d}{dx} G(x) \right) \frac{dx}{x} .$$

The function on the left hand side of (*) has poles at $s = 0, 1$ ([EdH] 1.8), whereby the pole at $s = 0$ is caused by the Gamma function. Therefore, Riemann multiplies it by $s(s-1)$ to define the entire function $Z(s)$; that is an analytical function, which is defined for all values of s , and the functional equation of the Zeta function is equivalent to $Z(s) = Z(1-s)$.

The functional equation is derived from the summation formula

$$\sum_{u \in \mathbb{Z}^n} e^{-t|u|^2} = \left(\frac{\pi}{t}\right)^{n/2} \sum_{u \in \mathbb{Z}^n} e^{-|\pi u|^2/t} , \quad t > 0 .$$

It is an example applied to the Gaussian function of the general Poisson summation formula ([PeB], Corollary 11.9)

$$\sum_{u \in \mathbb{Z}^n} \hat{\varphi}(2\pi u) = \sum_{u \in \mathbb{Z}^n} \varphi(u) \text{ and } \sum_{u \in \mathbb{Z}^n} \varphi(2\pi u) = (2\pi)^{-n} \sum_{u \in \mathbb{Z}^n} \hat{\varphi}(u) .$$

The functional equation is applied to derive the Riemann (error) function for the density function ([EdH] 1.13)

$$J(x) = \frac{1}{2\pi i} \int_{a-i\infty}^{a+i\infty} \log z(s) x^s \frac{ds}{s} .$$

The factor $(s-1)$ leads to the Li-function, while the factor $\frac{s}{2}$ is anticipated in the function $\gamma\left(1 + \frac{s}{2}\right)$ leading to the famous Riemann error function ([EdH] 1.16). Its convergence property jeopardizes a proof of the RH criterion

$$\pi(x) = Li(x) + O(\sqrt{x} \log x) = Li(x) + O(x^{\frac{1}{2}+\varepsilon}) , \quad \varepsilon > 0 .$$

Our proposed Kummer function based Zeta function theory is based on the Hilbert transform of the Gaussian function, which is the Dawson function

$$F(z) := e^{-z^2} \int_0^z e^{-t^2} dt \quad \text{resp.} \quad \sqrt{\pi} f_H(x) := F(\sqrt{\pi}x)$$

and its corresponding Mellin transform

$$M[f_H](s) = \pi^{(1-s)/2} \tan\left(\frac{\pi}{2}s\right) \gamma_{\frac{s}{2}}.$$

We note that $F(z)$ satisfies the differential equation $F'(z) + 2F(z) = 1$, leading to the (polynomial) asymptotic $F(\sqrt{x}) = O(x^{-\frac{1}{2}})$, alternatively to the asymptotic $f(\sqrt{x}) = O(e^{-x})$.

With respect to Riemann's "trick" above to build an entire function, this corresponds to an alternative multiplication with the factor $(s-1)\tan\left(\frac{\pi}{2}s\right)$ of (*).

We note the identity $\tan\left(\frac{\pi}{2}s\right) = \cot\left(\frac{\pi}{2}(1-s)\right)$; on the critical line $(s = \frac{1}{2} + it, t \in \mathbb{R})$ it holds $(\tan\left(\frac{\pi}{2}s\right) = \cot\left(\frac{\pi}{2}\bar{s}\right)$ (see also ([EdH] 10.2, 10.3, 10.10).

Related to the $\cot(\pi z)$ -function L. Kronecker provided some representation of series by integrals ([HeK] p. 42 ff):

via integration of the function $\pi \cot(\pi z) f(z)$ along of two parallel abscissa axis (with corresponding domain and convergence assumptions regarding the function $f(z)$) it holds the partial fraction expansion

$$\pi \cot(\pi z) f(z) = \sum_{k=-\infty}^{\infty} \frac{f(k)}{z-k}.$$

Also via integration along two parallel axis of ordinates there is a corresponding representation in a different form given by

$$\frac{1}{2} \left\{ \sum_{x_0 < k < x_1} f(k) + \sum_{x_0 \leq l \leq x_1} f(l) \right\} = -\frac{1}{2} \int_0^\infty (-1)^\alpha \pi \cot(x_\alpha + i\epsilon y) f(x_\alpha + i\epsilon y) dy$$

with $\alpha = 0, 1, \epsilon = -1, 1$ whereby $(x_0, x_1), (y_0, y_1)$ are the two abscissa resp. ordinates values with $(x_0 < x_1, y_0 < y_1)$.

From [EdH] 1.11 we recall the Stieltjes integral representation with the integrand of the log-Zeta function given by

$$\int_0^\infty x^{-z} dJ(x) = \frac{1}{2} \int_0^\infty x^{-z} d \left\{ \sum_{p^n < x} \frac{1}{n} + \sum_{p^l \leq x} \frac{1}{l} \right\}.$$

With respect to (***) above it formally also holds ([EdH] 10.3)

$$\frac{Z(s)}{s(s-1)} = \frac{1}{2} \int_0^\infty x^{1-s} G(x) \frac{dx}{x}$$

that is, the function is formally the transform of the operator

$$g(s) \rightarrow \int_0^\infty g(xs) G(x) dx.$$

But the above operator has no transform at all, as the integral does not converge for any s .

Nevertheless, this statement can be given substance as follows ([EdH] 10.5), which motivated our proposed alternative transform representation below:

A continuous analog of Euler's ludicrous formula

$$\sum_{n=-\infty}^{\infty} x^n = (1 + x + x^2 + \dots) + (x^{-1} + x^{-2} + \dots) = \frac{1}{1-x} + \frac{1}{x-1} = 0$$

is

$$\int_0^{\infty} x^{1-s} \frac{dx}{x} = \int_0^1 x^{1-s} \frac{dx}{x} + \int_1^{\infty} x^{1-s} \frac{dx}{x} = \frac{1}{1-s} + \frac{1}{s-1} = 0 .$$

This is, of course, nonsense because the values of s for which the above integrals converge are mutually exclusive – the first one being convergent for $\text{Re}(s) < 1$ and the second one being convergent for $\text{Re}(s) > 1$, but it does suggest that the formal transform of $g(x) \rightarrow \int_0^{\infty} g(ux)dx$ is zero.

The Hilbert transform is the proposed additional tool to overcome the above challenge to build a self-adjoint operator enabling the Berry-Keating (Hilbert-Polya) conjecture, as any Hilbert transformed function has vanishing constant Fourier term.

The Hilbert transform of the Gaussian function is the Dawson function with the asymptotic

$$f_H(\sqrt{x}) = O(x^{-\frac{1}{2}}) , x \rightarrow \infty .$$

Let $g(x) := e^{-x}$ and $I(x) = \int_{-\infty}^{\infty} \frac{g(t^2)}{x-t} dt$, then this leads to the replacement

$$\gamma(s) = M[g](s) \rightarrow M[I(\sqrt{x})](s) = \pi \cdot \tan(\pi s) \cdot \gamma(s) .$$

Let M and H denote the Mellin resp. the Hilbert transformation operators, then it holds

$$M \left[-x \frac{d}{dx} f(x) \right] (s) = sM[f](s) = \pi^{-\frac{s}{2}} \gamma \left(1 + \frac{s}{2} \right)$$

$$M \left[-\frac{d}{dx} (x^2 \frac{d}{dx} (f(x))) \right] (s) = (s-1)sM[f](s) = (s-1)\pi^{-\frac{s}{2}} \gamma \left(1 + \frac{s}{2} \right)$$

whereby ([EdH] 1.8 (1))

$$Z(s) = (s-1)\pi^{-\frac{s}{2}} \gamma \left(1 + \frac{s}{2} \right) Z(s) .$$

In order to anticipate the non-vanishing constant Fourier term of the Gaussian function, Riemann modified the above Mellin transform representation in the form ([EdH] 10.3)

$$Z^{**}(s) = \frac{1}{2} \int_0^{\infty} x^{1-s} \frac{d}{dx} (x^2 \frac{d}{dx} (G(x) - 1)) \frac{dx}{x}$$

which even destroys the (only formally valid) self-adjoint transform representation above (see also ([EdH] 12.5 for Polya's alternative, entire Zeta function definition linked to the Bessel functions).

With respect to the Hilbert-Polya conjecture this leads to a replacement of the above by

$$M \left[-\frac{d}{dx} (x f_H) \right] (s) = (s-1)sM[f_H](s) = c_1 \frac{\tan \frac{\pi s}{2}}{\frac{\pi}{2^s}} (s-1)\pi^{-\frac{s}{2}} \gamma \left(1 + \frac{s}{2} \right) ,$$

which then defines an alternative entire Zeta function in the form

$$Z^*(s) := c_1 \frac{\tan \frac{\pi s}{2}}{\frac{\pi}{2} s} Z(s) = c_1 \frac{\cot(\frac{\pi}{2}(1-s))}{\frac{\pi}{2} s} Z(s) = c_2 \left[-\frac{\pi}{2} (1-s) \cot\left(\frac{\pi}{2}(1-s)\right) \right] \pi^{-\frac{s}{2}} \gamma\left(\frac{s}{2}\right) Z(s)$$

and its corresponding transform representation given by

$$Z^*(s) = \frac{1}{2} \int_0^\infty x^{1-s} \frac{d}{dx} (x G_H(x)) \frac{dx}{x} .$$

The Gaussian function and its Hilbert transform are norm-equivalent with respect to the $L_2 = H_0$ –norm, i.e. both are equal in a weak L_2 –sense. The convergence of the transform representations in the critical stripe resp. on the critical line is ensured in a weak $H_{-\alpha}$ –sense ([EdH] 9.7, 9.8).

Both functions, $Z(s) Z^*(s)$ do have the same zeros. Therefore, in case the function $\theta^*(t) := Z^*\left(\frac{1}{2} + it\right)$ can be realized as a convolution $\theta(t) := (G * dF)(t)$ this would prove the RH ([CaD]). The proposed Hilbert space framework provides corresponding PDO with appropriate symbols, which goes in line with corresponding convolution integral operators.

The Berry conjecture is that the nonimaginary solutions E_n of the zeros $z_n = 1/2 + E_n$ of the Riemann Zeta function are the eigenvalues of an appropriate Hermitian operator H , providing a model for the quantum chaos [BeM]. The newly proposed Hilbert space Hilbert space for a quantum state Hilbert space framework is about a replacement of $L_2 \rightarrow H_{-1/2}$ with a corresponding replacement of the (energy space) domain of corresponding Schrödinger operator (in weak variation representation) $H_1 \rightarrow H_{1/2}$. The replacement

$$M \left[-\frac{d}{dx} (x^2 \frac{d}{dx} (f(x))) \right] (s) \rightarrow M \left[-\frac{d}{dx} (x f_H) \right] (s)$$

for the proposed RH solution goes along with this reduced regularity requirement. The leads to the *Calderón-Zygmund integral-differential operator* with symbol

$$|\omega| = \sum_{k=1}^n \omega_k \frac{\omega_k}{|\omega_k|}$$

defined on corresponding domain given by ([EsG] example 3.4, [LiI] example 3.1.4, 3.1.6)

$$L[u] := \sum_{k=1}^n Y_k D_k u = \sum_{k=1}^n \frac{\gamma\left(\frac{n+1}{2}\right)}{\pi^{\frac{n+1}{2}}} \text{p. v.} \int_{-\infty}^{\infty} \frac{x_k - y_k}{|x - y|^{n+1}} \frac{\partial u(y)}{\partial y_k} dy$$

resp

$$\Lambda^{-1} u = \frac{\Gamma\left(\frac{n-1}{2}\right)}{2\pi^{(n+1)/2}} \int_{-\infty}^{\infty} \frac{u(y) dy}{|x - y|^{n-1}}$$

where Y_k denote the Riesz operators, which are the n -dimensional generalizations of the Hilbert transform. The operator $L[\circ]$ is proposed alternatively to the “standard” differential operator to enable an alternative Schrödinger momentum operator and a corresponding alternative ground state energy model with non-integer Hilbert space domain ([BrK1], [BrK2]) and inner product definitions for differentials in the form ($u, v \in L_2$)

$$((du, dv)) := (L[du], L[dv])_{-1} , \quad ((du, v)) := (L[du], L[dv])_{-\frac{1}{2}} .$$

In Plemelj’s concept of an alternative simple and double layer potential with reduced regularity requirements to the underlying domains the corresponding differentials are named as “mass elements/ particles” in opposite to mass densities. The rotation invariance property of the Riesz operator ensures the “rotation invariance” of the considered differentials.

With respect to the entire Riemann Zeta function Z , the \cot – function and the Hermite polynomials h_n (with $\widehat{h}_n(\omega) = (-i)^n h_n$) we further note that $Z \in H_{-1}$, $\cot \in H_{-1}^\#$, $h_n \in L_2$, i.e. $(Z, \log \sin)_{-1/2}$, $(Z - \cot, \log \sin)_{-1/2}$, $(Z, h_n)_{-1/2}$, $(Z - \cot, h_n)_{-1/2}$ are defined. In this context we refer to

- the Bagchi reformulation of the Beurling RH criterion, as the $H_{-1/2}$ – Hilbert space is dense in H_{-1} with respect to the $\| \cdot \|_{-1}$ norm
- the wavelet section below
- the theory of cardinal series ([WhJ] §11).

The Calderón-Zygmund integral-differential operator is also proposed as alternative Schrödinger (-Calderón) momentum operator with domain $H_{1/2} = H_0 + H_0^-$. The corresponding *inner product* definition $((du, dv))$ is provided in [BrK1]. This alternative Schrödinger momentum operator definition provides additional evidence to the famous constant $-\frac{1}{i} = i$ of the Schrödinger momentum operator in its classical form. For space dimension $m = 1$ it is given by $u(x) \rightarrow P^*[u](x) := -i \frac{d}{dx} H[u](x) = -iH[u_x](x)$ ([MeY], 7.1). With respect to the $H_{-1/2}$ – inner product it holds $(P^*[u], v)_{-1/2} \cong (u, v)_0$. In the simplest case (1-D periodical functions) it is given by the convolution (Stieltjes/Plemelj) integral

$$\int \cot\left(\frac{x-y}{2}\right) du_y.$$

In the context of the newly proposed “energy-space” $H_{1/2} = H_1 + H_1^-$ we further refer to the Bose-Einstein condensation, where below the critical temperature T_c BEC “normal gas” particles coexist in equilibrium with “condensed” particles. Unlike a liquid droplet in a gas, here the “condensed” particles are not separated in space ($H_{-1/2} = H_0 + H_0^-$) from normal particles. Instead they are separated in momentum space. The condensed particles all occupy a single quantum state of zero momentum, while normal particles all have finite momentum.

In case the domain of such a compact operator is the L_2 Hilbert space the corresponding eigenfunctions build the basis of this Hilbert space. The concept of “wave package” enables also continuous spectra. Therefore, such “wave packages” require a domain extension (e.g. $L_2 \rightarrow H_{-1/2}$) in order to ensure convergent inner products and related norms. “Wave packages” are also called “eigen-differentials” (H. Weyl), playing a key role in quantum mechanics in the context of the spectral representation of Hermitian operators (D. Hilbert, J. von Neumann, P. A. M. Dirac).

We note the Gaussian function related “Mexican hat” function

$$g(x) := -\frac{d^2}{dx^2} \left(e^{-\frac{x^2}{2}} \right) = (1 - x^2) e^{-\frac{x^2}{2}}.$$

being successfully applied e.g. in wavelet theory (see below).

In [StA] *variational space-time integrators for Maxwell's equations* are developed. The concept combines spatial and time discretization in developing *geometric numerical integrators*. The approach preserve, by construction, various geometric properties and invariants of the continuous physical systems that they approximate. With respect to "the nonlinear Landau damping for general interaction" we note that the *variational space-time integrators* concept treats electromagnetic Lagrange density as a discrete differential 4-form in space-time, while in [MoC], [ViC] the analysis is concerned

- with Fourier analysis with respect to the space dimension in combination with C^0/L_∞ –estimates with respect to the time dimension. It ends up in "analytical norm" estimates requiring mathematical regularity assumptions, which are not appropriate for the underlying physical (plasma) models (e.g. KAM theory).
- with different regularity requirements concerning the most critical underlying interaction potential W ($W \in H_{-1/2-\varepsilon}$, $W \in H_{-1}$) of the Vlasov-equation.

The Landau equation (a model describing time evolution of the distribution function of plasma consisting of charged particles with long-range interaction) is about the Boltzmann equation with a corresponding Boltzmann collision operator where almost all collisions are grazing (based on a Fourier multiplier representation ([LiP1]) in line with Oseen kernels, Laplace and Fourier analysis techniques [LeN] and scattering problem analysis techniques based on Garding type (energy norm) inequalities (like the Korn inequality)). *Its solutions enjoy a rather striking compactness property.*

The collision operator is given by

$$Q(f, f) = \frac{\partial}{\partial v_i} \left\{ \int_{\mathbb{R}^N} a_{ij}(v-w) \left[f(w) \frac{\partial f(v)}{\partial v_j} - f(v) \frac{\partial f(w)}{\partial w_j} \right] dw \right\}$$

with

$$a_{ij}(z) = \frac{a(z)}{|z|} \left\{ \delta_{ij} - \frac{z_i z_j}{|z|^2} \right\} = \frac{a(z)}{|z|} P(z) := \frac{1-[1-a(z)]}{|z|} [Id - Q](z) \quad Q(z) := (R_i R_j)_{1 \leq i, j \leq N}$$

and $a(z)$ symmetric, non-negative and even in z and R_i denote the Riesz operators.

Our 3D-Navier-Stokes equations (NSE) solution is based on a distributional Hilbert space concept to derive appropriate energy norm estimates of the non-linear, non-stationary 3-D-Navier-Stokes equations. In [BrK3] we propose problem adequate norms to derive corresponding adequate a priori estimates for the transport equation, which are in line with the results of [LiP], [LiP1].

The variation theory approach in appropriately defined Hilbert spaces is proposed to prove the Landau damping based on the Landau equation. A weak variational representation (which is anyway the proper framework for quantum theory) would overcome Vlasov's mathematical argument against the Landau equation that "this model of pair collisions is formally (!) not applicable to Coulomb interaction due to the divergence of the kinetic terms". At the same point in time the "analytical norm" proof of the Landau damping effect ([MoC]) based on the Vlasov equation does not reflect the underlying physical explanation of this effect. This is about "wave damping w/o energy dissipation by collisions in plasma", because electrons are faster or slower than the wave and a Maxwellian distribution has a higher number of slower than faster electrons as the wave. As a consequence, there are more particles taking energy from the wave than vice versa, while the wave is damped ([BiJ]).

A "helicopter view" regarding the "Higgs field" and "Kant's ether field"

The gauge invariance is the main principle in current SMEP theory.

[BID] 10.3: *"It is fine that the gauge field of electromagnetism has zero mass because there the force is mediated by photons, which are massless. However, Yang-Mills type forces must arise from the exchange of massive particles because of the observed short range of these forces. The Higgs mechanism helps in two ways. First, gauge fields can acquire mass by the symmetry breaking. Second, the undesirable Goldstone bosons (which arise in the symmetry-breaking process) can be usually gauged away."*

The Higgs effect (or mechanism) builds on an extended from global to local $U(1)$ transformations symmetry group of the underlying Lagrangian. It explains the mass of the gauge W- and Z-(weak interaction) bosons of the weak "nuclear-force".

[HiP]: *"Within the framework of quantum field theory a "spontaneous" breakdown of symmetry occurs if a Lagrangian, fully invariant under the internal symmetry group, has such a structure that physical vacuum is a member of a set of (physically equivalent) states which transform according to a nontrivial representation of the group. This degeneracy of the vacuum permits non-trivial multiplets of scalar fields to have nonzero vacuum expectation values (or "vacuons"), whose appearance leads to symmetry-breaking terms in propagators and vertices. ... When the symmetry group of the Lagrangian is extended from global to local transformations by introduction of coupling with a vector gauge field the original scalar massless boson as a result of spontaneous breakdown of symmetry then becomes the longitudinal state of a massive vector (Higgs) boson whose transverse state sare the quanta of the transverse gauge field. A perturbative treatment of the model is developed in which the major features of these phenomena are present in zero order."*

The Higgs boson is supposed to be a heavy elementary particle (with non-zero rest mass of about 125 GeV with spin 0). The Higgs field is supposed to fill the whole universe interacting with each particle, which "moves" through it by a kind of frictional resistance, i.e. which has kinetic energy. Therefore, the Higgs effect (i.e. generating mass particles) requires a Higgs field with not vanishing amplitudes in the ground state. The corresponding Lagrange density is described by the (Lorentz-invariant) Klein-Gordon equation (i.e. it is about a relativistic theory) equipped with the Higgs potential, which is a "special" "Mexican hat" potential in the form

$$V(\varphi) := \mu^2 \varphi \varphi^* + \beta (\varphi \varphi^*)^2.$$

The corresponding local gauge invariant quantum field theory is about the related Lagrange density, which is invariant under a local $U(1)$ gauge transformation, i.e. there is an infinitesimal symmetry transformation of matter & gauge fields (fermions & bosons). The Higgs effect overcomes the issue of the weak nuclear power SM model that there can be existed no particles with mass w/o violating the gauge invariance of the describing Lagrange density.

The current understanding of all known "particles" in the universe is, that there is a split into two groups of those "particles" to overcome the contact body (body-force interaction) problem

1. spin(1/2)-"matter"- "particles", which are "objects" with a spin(1/2), i.e. those "objects" *look the same* only after the second rotation
2. spin(0,1,2)-"force"- "particles", which are "objects" with spins 0,1,2, interacting with spin(1/2)-"matter" "objects".

The first group goes back to Dirac, who introduced this *purely mathematical* concept to "explain" why spin(1/2)-"matter"- "particles", especially the electrons, can exist as "separate" "objects", while not merging to one big "soup" ("object"?). Dirac's theory enables consistency of the quantum mechanics and the special relativity theory.

In order to avoid the same ("soup" disaster) effect Pauli postulated his exclusion principle in order to ensure that spin(1/2)-"matter"- "particles" under the influence of spin(0,1,2)- "force"- "particles" do not collapse to a state of extremely high density.

We mention the following open challenges:

- *the Higgs effect still leads to oppositions for energies not in the range of $7\text{GeV} < m_H < 1\text{TeV}$*
- *the gauge invariance principle based on the Lagrange formalism leads to bosons with mass to generate fermions, while the original concept of fermions and bosons was to distinguish between particle with mass (fermions) and mass-less "force" interaction particles between them (bosons)*
- *a "gravitation" "boson" is not possible to be adapted to the SMEP, while with the Higgs boson there is now a "particle" included, which "acts" in the whole universe, especially in the vacuum (which fills most of the space, in the cosmology world, but also in the quantum world), but only affecting weak nuclear-forces and its related W- and Z-bosons.*

The alternative approach of this paper is about a replacement of the gauge invariance principle per defined (problem specific) Lagrange density by a "rotation invariance" principle per defined Hilbert space framework (which is common for all physical problem specific Lagrange density representations). The beauty of the gauge methodology to gauge away undesirable (e.g. Goldstone) bosons has its counterpart in the Einstein field equation by the covariant derivative concept going in line with *differentiable* manifolds (whereby the differentiability requirement is w/o any physical meaning). The corresponding concept, which replaces the inertial system concept of the Newton gravity model, is an infinitesimal displacement (tensor) fields A_j^k with $dA^k - \delta A^k = A_j^k dx^j$. Our alternative approach avoids the above purely mathematical model driven concepts, which are w/o any physical meaning.

As a consequence, the proposed perturbative treatment of the Higgs model ([HiP]) is replaced by a corresponding approximation analysis of compactly embedded closed spaces in the considered Hilbert space framework (see e.g. ([AzA], [CoD])).

The mass of a proton consists nearly exclusively of the energy of the gluons, which are the interconnection particles, which hold together the quark. In this sense, mass is essentially the manifestation of the vacuum energy (which could be interpreted as a projection operator from a "vacuum" field into a "matter field"). The "Higgs" field, interpreted as a kind of ether which exists in the whole universe, goes along with "Kant's conception of Ether as a field" ([WoW]), and his ontology of space, which is anti-Newtonian in the sense that space is not an object, but the form of representing a physical object. As there is only about 10% of matter of the universe "covered" by (obviously inconsistent) mathematical physics models (e.g. quantum and gravitation field theory) while about 90% of the remaining matter is unknown "dark" matter with not-known form and substance "embedded (?)" in a "universe space", which mainly "contains" "vacuum" besides the above 100% matter it is might be worth to review Kant's dynamic view of space ([FöE]).

With respect to the alternative proposed "quantum state" Hilbert space model of this paper (where the "Higgs" particles might be just another name of those quantum particles) we note that it enables

- a projection operator $P: H_{-\alpha} \rightarrow H_0$ to model "matter creation" from purely mass-less generalized "wave/energy" quantum states out of the complementary closed space $H_{-\alpha} - H_0$
- the Hamiltonian formalism based on the variational "energy" representation of appropriate PDO equations with reduced regularity requirements compared to the Lagrange formalism; in those cases the Legendre ("contact" (!)) transformation cannot be applied to prove the equivalence of both formalisms
- an infinitesimal *rotation invariant* symmetry (invariance), based on a variational (weak) Hamiltonian formalism, i.e. any "renormalization" requirements based on local, (only) affine infinitesimal transformations are not required anymore.

The Maxwell equations: baseline for the Yang-Mills and the gravity field equations

The Maxwell equations (and its related (complementary) variational principles with its underlying $U(1)$ symmetry group) build the baseline for all related SMEP Lagrange density formulations and the Einstein field equations:

[EiA] p. 52 ff: „die Maxwellschen Gleichungen bestimmen das elektrische Feld, wenn die Verteilung der elektrischen Ladungen und der Ströme bekannt sind. Die Gesetze aber, nach denen sich Ströme und Ladungen verhalten, sind uns nicht bekannt. Wir wissen wohl, dass die Elektrizitäten in Elementarkörperchen (Elektronen, positiven Kernen) bestehen, aber wir begreifen es nicht vom theoretischen Standpunkt aus. ... Wir kennen daher, falls wir überhaupt die Maxwellschen Gleichungen zugrunde legen dürfen, den Energietensor für die elektromagnetischen Felder nur außerhalb der Elementarteilchen. Wir wissen heute, daß die Materie aus elektrischen Elementarteilchen aufgebaut ist, sind aber nicht im Besitz der Feldgesetze, auf welchen die Konstitution jener Elementarteilchen beruht. Wir sind daher genötigt, uns bei deren Behandlung der mechanischen Probleme einer ungenauen Beschreibung der Materie zu bedienen, welche der von der klassischen Mechanik verwendeten entspricht. Die Dichte σ der ponderablen Substanz und die hydrodynamischen Druckkräfte (Flächenkräfte) sind die Grundbegriffe, auf die eine derartige Beschreibung sich stützt“.

In classical electromagnetic field theory one deals with the following quantities: \mathbf{E} = electric field, \mathbf{H} = magnetic field, \mathbf{B} = magnetic induction, \mathbf{J} = electric current density, \mathbf{D} = dielectric displacement ρ = charge density. In the language of exterior forms this leads to ([FIH], 4.6)

$$d\alpha = 0 \quad , \quad d\beta = -4\pi\gamma \quad (\alpha = \alpha_{E,B}, \beta = \beta_{H,D})$$

while $d\gamma = 0$ corresponds to $\dot{\rho} + \text{div}J = 0$ ($\gamma = \gamma_{J,\rho}$). Maxwell equations in free spaces are simply

$$d\alpha = 0 \quad d*\alpha = 0 \quad .$$

With respect to the “rotation” invariance topic of this paper we note the integrated Maxwell field equations by Lorentz ([WeH] §20) based on the assumption that the distribution of charges and currents are known:

The equation $\text{div}\mathbf{B} = 0$ is satisfied by setting $-\mathbf{B} = \text{curl}\mathbf{f}$ in which \mathbf{f} is the *vector potential*. By substituting this in the first Maxwell equation $\text{curl}\mathbf{E} + \frac{1}{c}\frac{\partial\mathbf{B}}{\partial t} = 0$ one gets that $\mathbf{E} - \frac{1}{c}\frac{d\mathbf{f}}{dt} = 0$ is irrotational so that one can set $\mathbf{E} - \frac{1}{c}\frac{d\mathbf{f}}{dt} = \text{grad}\varphi$ in which φ is the *scalar potential*. Finally the vector and scalar potential are defined by correspond wave equations in the following form (based on the differential form representation of the Maxwell equations)

$$\frac{1}{c^2}\ddot{\varphi} - \Delta\varphi = -\rho \quad , \quad \frac{1}{c^2}\ddot{\mathbf{f}} - \Delta\mathbf{f} = -\mathbf{s} \quad .$$

We note that the Klein-Gordon equation is the relativistic counterpart of the wave equation. The Maxwell equations reflect Faraday’s near action theory approach. They are based on Gauss’s law for electric fields, which are formulated in integral and differential form requiring different regularity assumptions for the different domains of the corresponding operators. Therefore, the first step to enable a “rotation invariance” principle based on common domains for all SMEP affected operators, is to consider only the integral form of the Gauss’s electric fields, i.e.

$$\oint_S \vec{E} \circ \vec{n} da = \frac{q_{encl}}{\epsilon_0}$$

with the electric field \vec{E} , the permittivity of free space (or "vacuum permittivity") ε_0 , the amount of the enclosed contributing charge q_{encl} and Maxwell's displacement current

$$I_d := \varepsilon_0 \frac{d}{dt} \oint_S \vec{E} \circ \vec{n} da.$$

The second (main) step is to apply Plemelj's alternative definition of simple and double layer potentials (to further reduce regularity requirements of the affected operator domains), which goes along with Plemelj's revisited definitions of "mass & flux" vs. "mass densities" (see also the problem B related homepage: www.navier-stokes-equations.com). Plemelj's key concept (for space dimension $n = 2$) replaces the normal derivative concept by the integral (σ, σ_0 elements of the boundary!)

$$\bar{U}(\sigma) = - \int_{\sigma_0}^{\sigma} \frac{dU}{dn} ds,$$

whereby $d\bar{U}(\sigma)$ may be defined, but not $\frac{dU}{dn} = - \frac{d\bar{U}(\sigma)}{d\sigma}$. The new term is named "current strength" flowing through the corresponding boundary piece, also called "current or flux".

In case of the logarithmic potential (for space dimension $n = 2$) the current is determined by a simple measure: it is the change of the conjugate potential (which is related to the Hilbert transform) between the points σ_0, σ . The generalization to the space dimensions $n > 2$ is provided in [StE] based on the Riesz operator concept. With respect of the relationship of the YME and the Hilbert transform we refer to [DuR].

Plemelj's quotes [PIJ]: §5: "es handelt sich hier um eine Verallgemeinerung, wie es die Erweiterung differentiierbarer Funktionen auf die stetigen ist". §8: „Bisher war es üblich für das Potential $V(p)$ die form

$$\int \log \frac{1}{r_{ps}} \dot{\mu}(s) ds$$

vorauszusetzen, wobei dann $\dot{\mu}(s)$ die Massendichtigkeit der Belegung genannt wurde. Eine solche Annahme erweist sich aber als eine derart folgenschwere Einschränkung, dass dadurch dem Potentiale der größte Teil seiner Leistungsfähigkeit hinweg genommen wird. Für tiefergehende Untersuchungen erweist sich das Potential nur in der (Stieltjes-schen) Form

$$\int \log \frac{1}{r_{ps}} d\mu_s$$

verwendbar.

It enables a replacement of the

- "Dichte (density) $\sigma = \mu(s)ds$ der ponderablen Substanz" ([EiA] p. 52) by the Plemelj "mass element $d\mu_s$ "
- "hydrodynamischen Druckkräfte (Flächenkräfte (area forces))" ([EiA] p. 52) by the Plemelj „current“
- co-variant (Maxwell electromagnetic *potential*) vector by the Plemelj "potential".

In other words, Plemelj's (Stieltjes integral based) mass element and potential definitions transform Newton's (Lebesgue integral based) mass density $\mu(s)ds$ and potential definition moving from a far distance action theory to a near action theory. In combination with a correspondingly defined (rotation invariant) inner product for those "mass elements" an adapted variational representation of the Maxwell equations is proposed as model to describe the behavior of currents and charges of an electromagnetic field. We note that the co-variant derivative of a scalar field corresponds to the partial derivative.

In section G. below we briefly sketch the relationship of Plemelj's alternative potential definition, the Hilbert transform and the Laplacian equation, which may be solved using simple layer potential. The 3-D analogy of the Cauchy-Riemann differential equations is provide in [RuC]. The Riesz transforms on spheres are provide in [ArN].

The achieved regularity reduction, while the Green formula keep still valid, enables a redefined Dirichlet (energy) integral $D(u, v) := (\nabla u, \nabla v)$ in the form ($\alpha = 0$ being replaced by $\alpha = -1/2$)

$$(u, \Delta v)_\alpha + (\nabla u, \nabla v)_\alpha = (u, \Delta v)_\alpha + (u, v)_{1*\alpha} = \langle u, v_n \rangle_\alpha .$$

With respect to the Lorentz integration of the Maxwell equations above this goes along with a variational representation in case of the first wave equation (which denotes a wave disturbance travelling with velocity c) in the form

$$\varphi \in H_{1/2} : \frac{1}{c^2} (\ddot{\varphi}, \dot{\vartheta})_{-1/2} + (\varphi, \vartheta)_{1/2} + \langle \varphi_n, \vartheta \rangle_{-1/2} = -(\rho, \vartheta)_{-1/2} \quad \forall \vartheta \in H_{1/2}$$

The wave equation plays also a key role in the harmonic quantum oscillator analysis. The variational representation is basically about hyper-singular integral equations. In [WeP], [WeP1] the Laplace operator with respect to electric and magnetic boundary conditions is analyzed. In [ChF] a method is introduced and analyzed allowing appropriate approximations w/o the necessity of a Lagrange multiplier.

The corresponding Klein-Gordon equation $\left[\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \Delta + \frac{m^2 c^2}{\hbar^2} \right] \varphi(x) = 0$ is derived from the energy-momentum equality

$$(*) \quad E^2 - \mathbf{P}^2 c^2 = m^2 c^4 I$$

whereby $E := i\hbar \frac{\partial}{\partial t}$ and $\mathbf{P} := -i\hbar \nabla$ denotes the standard Schrödinger operator. Dealing with the quadratic terms of (*) leads to also negative energy values E ([HeW] (134)).

Replacing the standard Schrödinger operator in (*) by the proposed one of this paper, leads to $\mathbf{P}^{*2} = -\hbar H^2 \left[\frac{d^2}{dx^2} \right] = \hbar \frac{d^2}{dx^2}$ (because of $H^2 = -I$) and therefore, to

$$(**) \quad \frac{1}{c^2} E^2 + \mathbf{P}^{*2} = m^2 c^2 I$$

which indicates a Helmholtz type equation given by

$$-\nabla^2 u - \mu^2 u = \left\{ \begin{array}{l} \delta_i \\ 0 \end{array} \right. \text{ in the considered domain}$$

(δ_i denotes the Dirac function at source i corresponding to the fundamental solutions vs. zero for the general solution) with its underlying distance variable (Hankel function based) solutions governed by the divergence (conservation) theorem and the Sommerfeld radiation condition at infinity ([KyP]). It enables the so-called distance function wavelet analysis based on Helmholtz-Fourier transforms and corresponding Helmholtz-Fourier series (e.g. [AmS], [ChW]). For its related Riesz transforms on spheres we refer to [ArN].

We mention that the Helmholtz (decomposition) theorem states that any sufficiently smooth, rapidly decaying vector field in three dimensions can be resolved into the sum of an irrotational (curl-free) vector field and a solenoidal (divergence-free) vector field.

A "helicopter view" regarding "differentials" and "distributional Hilbert scale"

The current quantum state Hilbert space is identical with the "measurement /observation /statistical" function space L_2 . The newly proposed distributional Hilbert spaces are $H_{-1/2}$ and H_{-1} . The corresponding inner product $((x,y)_{-1}$ can be put into relation to an inner product in the form $((dx,dy))$. The role of the L_2 as the quantum state Hilbert space is being replaced by $H_{-1/2}$.

As the L_2 is a closed, compactly embedded subspace of $H_{-1/2}$ with respect to the $H_{-1/2}$ -norm, resp. as the $H_{-1/2}$ is a closed, compactly embedded subspace of H_{-1} with respect to the H_{-1} -norm, compactness arguments can be applied in combination with corresponding Garding type inequalities of related operator representation in the form $B = A + K$ ($[AzA]$, $[BrK]$).

The differential "objects" can be interpreted as "ideal point" (non-standard numbers, "monads"), which then are going to replace the real numbers. Both fields do have the same cardinality, but the non-standard numbers allow infinitesimal small number "objects" in the neighborhood of each "real" number. In a certain sense the field of real numbers is compactly embedded into the field of non-standard numbers. The Riesz theorem below provides the corresponding relationship to a closed subspace X_0 of a Hilbert space X .

The advantages with respect to the three considered problems are the following

1. the "measurement of real numbers is already an approximation by rational numbers, i.e. truly "observations" of irrational number "objects" are not possible; each irrational number is already a full universe, i.e. an approximation of an infinite numbers of rational numbers; extending those number field to ideal numbers is just the same mystery with same cardinality; the key differentiator is related to a measurement of length axiom by given "unit of measure" length

2. the $H_{-1/2}$ Hilbert space provides an alternative model to the current $H_0 = L_2$ "quantum state" Hilbert space, which avoids the Dirac "function" concept with its handicap of space dimension depending regularity requirements ($\delta \in H_{-\frac{n}{2}-\epsilon}, \epsilon > 0$)

3. The physical "observation" L_2 Hilbert space (supporting also statistical analysis) is still valid and applicable; it is a subspace of $H_{-1/2}$, whereby its corresponding complementary closed subspace H_0^- of $H_{-1/2} = H_0 + H_0^- = L_2 + L_2^-$ enables an alternative modelling of "wave packages" ("eigen-differentials")

4. the Zeta function Z on the critical line is an element of the H_{-1} Hilbert space. Therefore, the proposed framework enables the Bagchi-Beurling (density) criterion, as both, the fractional part function $\rho(x)$ and the $\log(\sin(x))$ function, are elements of the Hilbert space L_2 , i.e. the inner products $(Z, \rho)_{-1/2}$, $(Z, \log(\sin))_{-1/2}$ are defined, and as the Hilbert space $H_{-1/2}$ is dense in H_{-1}

5. the $H_{-1/2}$ Hilbert ("fluid") space enables the Sobolevskii-estimate of the non-linear term of the 3-dimensional, non-stationary, non-linear variation NSE representation, leading to the bounded, generalized energy inequality

$$\frac{1}{2} \frac{d}{dt} \|u\|_{-1/2}^2 + \|u\|_{1/2}^2 \leq |(Bu, u)_{-1/2}| \leq c \cdot \|u\|_{-1/2} \|u\|_1^2$$

for $u \in H_1$.

The theorem of Riesz ensures "quasi-optimal" approximation properties of each "object" of a closed subspace X_0 of a Hilbert space X :

Theorem (Riesz): For each ε with $0 < \varepsilon < 1$ there exists a $y = x_\varepsilon \in X$ with $\|y\| = 1$ and

$$\inf\{\|x - y\| \mid x \in X_0\} \geq \varepsilon$$

Remark 1: A separable Hilbert scale can be built from the solutions of an eigenvalue equation

$$Kx = \sigma x$$

where K denotes a symmetric and compact operator:

Lemma: for no more than countable values σ_i the equation $Kx = \sigma x$ possesses non-trivial solutions x_i and $\lim_{i \rightarrow \infty} \sigma_i = 0$.

Remark 2: as 0 is an element of X_0 this means that the inf-term above is at most equal 1; therefore the theorem states that this value can be arbitrarily close approximated. This can be interpreted as counterpart of the approximation of an irrational number by rational numbers.

Remark 3: Regarding "ideal points in space-time" we refer to [GeR], where ... "a prescription is given for attaching to a space-time M , subject only to a causality condition, a collection of additional "ideal points". In particular, for any asymptotically simple space-times, the ideal points can be interpreted as the boundary at conformal infinity. The concept makes possible an extension of the domain-of-dependence concept to causal spaces. ... Some of these ideal points can be interpreted as singular points of M , others as points of infinity." In [PeR] the concept of a "conformal cyclic cosmology (CCC)" in combination with a revisited second principle of thermodynamics is proposed.

Theorem: Let H_1 and H_0 be Hilbert spaces (H_1 being a subspace of H_0) with

- i) $\|x\|_0 \leq \|x\|_1$ for $x \in H_1$
- ii) H_1 is dense in H_0
- iii) the unit ball of H_1 is relatively compact in H_0 .

Then there exists an operator A with $D(A) = H_1$, $R(A) = H_0$ and $\|x\|_1 = \|Ax\|_0$, whereby the operator A is

- i) positive definite, self-adjoint
- ii) A^{-1} is compact.

The corresponding eigenvalue problem

$$A\varphi_i = \sigma_i \varphi_i$$

has infinite solutions $\{\sigma_i, \varphi_i\}$ with $\sigma_i \rightarrow \infty$ and $(\varphi_i, \varphi_k) = \delta_{i,k}$, and for each element $x \in H_1 = A^{-1}H_0$ it holds the representation

$$x = \sum_{i=1}^{\infty} (x, \varphi_i) \varphi_i$$

Inner products with corresponding norms of a distributional Hilbert scale can be defined based on the eigen-pairs of an appropriately defined operator in the form

$$(x, y)_\alpha := \sum_i^\infty \lambda_i^\alpha (x, \varphi_i)(y, \varphi_i) = \sum_i^\infty \lambda_i^\alpha x_i y_i$$

Additionally, for $t > 0$ there can be an inner product resp. norm defined for an additional governing Hilbert space with an "exponential decay" behavior in the form

$$e^{-\sqrt{\lambda_i} t}$$

given by

$$(x, y)_{(t)}^2 := \sum_{i=1}^\infty e^{-\sqrt{\lambda_i} t} (x, \varphi_i)(y, \varphi_i)$$

$$\|x\|_{(t)}^2 := (x, x)_{(t)}^2$$

The distributional $H_{-1/2}$ – Hilbert space is proposed to model quantum states, alternatively to the Hilbert space H_0 . A mathematical (wavelet microscopic) analysis of those states is then about an analysis of the "objects"

$$x = x_0 + x_0^\perp \in H_0 \otimes H_0^\perp$$

with

$$\|x_0\|_0 = 1 \quad \sigma := \|x_0^\perp\|_{-1/2}^2$$

As it holds for any $t, \delta, \alpha > 0$ and $\lambda \geq 1$ the inequality

$$\lambda^{-\alpha} \leq \delta^{2\alpha} + e^{t(\delta^{-1} - \sqrt{\lambda})}$$

the following inequality is valid for any $x \in H_0$, governing the approximation "quality" of a quantum state with respect to the norm of H_0 :

$$\|x\|_{-1/2}^2 \leq \delta \|x\|_0^2 + e^{t/\delta} \|x\|_{(t)}^2 = \sigma \|x\|_0^2 + e \|x\|_{(\sigma)}^2 = \sigma \|x\|_0^2 + \sum_{i=1}^\infty e^{1-\sqrt{\lambda_i} \sigma} x_i^2$$

It balances the "continuous" view of the overall state with its "discrete" components.

A "helicopter view" on the unknown mathematical model of a "big bang" conform initial or radiation PDO equation model

Regarding the "big bang" singularity for $t \rightarrow 0$ and the newly proposed Hilbert space framework $H_{-1/2}$ we note that already the solution of the initial value problem of the (simplest, purely parabolic, linear) heat equation diverges for $t \rightarrow 0$ in case the initial value function is measurable, i.e. $u_0 \in H_0 = L_2$ ($\|u\|_k \leq ct^{-k/2}\|u_0\|$), while with reduced regularity assumption ($u_0 \in H_{-1/2}$) it holds $\|u\|_k \leq c\|u_0\|_{-1/2}$. The corresponding situation is valid for the non-stationary Navier-Stokes equations and the (free boundary) Stefan problem.

Beside the (well established) theory of initial value (parabolic) PDE problems there is the still open "Courant-Hilbert" conjecture concerning the most simple (hyperbolic, linear) initial value wave equation. The Huygens principle is valid under the same prerequisites for the initial value problem, as well as for the corresponding radiation problem ([CoR] VI, §9.1, §10.3). The "Courant-Hilbert conjecture" is about the reverse, i.e. distortion-free, progressive, spherical waves do only exist, if the Huygens principle is valid. In combination with the Hadamard conjecture ([CoR] §9.1) this would characterize the 4-dimensional space-time continuum and its related Maxwell theory.

Having in mind that already each irrational number is its own mathematical universe (i.e. it is defined by a sequence of an infinite numbers of rational numbers) the solution of the problem above seems to be a necessarily question to be answered, before a mathematical model of the "big bang" "situation" can be formulated.

In [PeR] a revisited entropy concept (see also "D" below) is proposed leading to a proposed conformal cycle cosmology, which is basically a periodical solution of a corresponding mathematical model, *that the universe as a whole is to be seen as an extended conformal manifold consisting of a (possible infinite) succession of aeons, and each appearing to be an entire expanding universe history* ([PeR]p. 147):

"According to current particle-physics about how to masses of basic particles actually come about, a particle's rest-mass ought to arise through the agency of a special particle (or perhaps a family of such special particles) referred to as a Higgs boson(s). ... In the very early universe, when the temperature (=energy form !!) was so high as to have provided energies greatly in excess of this Higgs value, all particles would then, according to standard ideas, indeed become effectively massless, like a photon."

The physical field, of which photons provide the quantum constituents, is the Maxwell electromagnetic field. It is completely conformal invariant.

In [FrH] the "hyperboloidal initial value" problem for Einstein's conformal vacuum field equations (reduced to Cauchy problems for first order quasilinear symmetric hyperbolic systems) is considered. For initial data of the class $H_s, s \geq 4$, there is a unique development which is a solution of the conformal vacuum field equations of class $H_s, s \geq 4$. This provides a solution of Einstein's vacuum field equations which has a smooth structure at past null infinity (note that according to the Sobolev embedding theorem it holds that the Hilbert space H_s is a subset of C^0 for $s > n/2$). The regularity of this solution is still a strong PDE solution, i.e. the correspondingly defined weak solution would even allow even in the standard case (w/o the proposed alternative energy space framework) regularity assumption of the class $H_s, s \geq 2$, which is still more regular than in the above considered parabolic initial value problems (heat equation and Stefan problem, as well as the NSE).

In [TaN] the theory of Cauchy problems for solutions of elliptic equations is provided covering PDO in the space of distributions on closed sets, generalized form of capacity associated with a semi normed space applied to systems of differential equations with injective (surjective) symbols.

The analogue concept of the Riemann curvature tensor in elasticity theory is the fourth-rank (elasticity) tensor, which links the stress tensor to the strain tensor by Hook's law $\sigma_{jk} = c_{jkmn}\varepsilon_{mn}$.

The first boundary value problem of elasticity is given by ([FiG])

given $\underline{f} \in \underline{L}_2$ find $\underline{u} \in \underline{H}_2$ such that

$$-\nabla\sigma(\underline{u}) = \underline{f} ; \quad -(\sigma_{ik,k}(\underline{u})) = f_i$$

It holds the shift theorem: for $\underline{f} \in \underline{L}_2$ the solution $\underline{u} \in \underline{H}_2$ exists uniquely and $\|\underline{u}\|_{H_2} \leq c \|\underline{f}\|_{L_2}$, i.e. the operator $-\nabla(\sigma(\underline{u}))$ behaves like the Laplacian operator. The corresponding weak variational representation of the boundary value problem is given by (Nitsche's method for contact problems)

$$a(\underline{u}, \underline{w}) = (\underline{f}, \underline{w}) , \quad \underline{w} \in H_1$$

with

$$a(\underline{v}, \underline{w}) := (\sigma_{ik}(\underline{v}), \varepsilon_{ik}(\underline{w})) + \oint n_i \{ \sigma_{ik}(\underline{v}) w_k + \sigma_{ik}(\underline{w}) v_k \} ds .$$

In [ShI] necessary and sufficient solvability conditions of ill-posed non-homogeneous Cauchy problems for PDE (as the Cauchy problem for the Lamé operator) with injective symbol of order ≥ 1 are provided.

In [CiP] an approach to the purely traction problem of the 3-dimensional linearized elasticity problem is provided, whose novelty consists in considering the linearized strain tensor as the "primary" unknown, instead of the displacement itself as is customary. The approach leads to a well-posed minimization problem, constrained by a weak form of the St Venant compatibility conditions. The corresponding complementary extremal problem (as alternative to the minimization problem of the potential energy) is based on the principle of Castigliano leading to the quadratic form $W(\varepsilon) = \frac{1}{2} c_{jkmn} \varepsilon_{jk} \varepsilon_{mn}$ ([VeW] 4.2.6).

With regards to the Maxwell equations we note that the components of the electric and magnetic field forces \underline{E} , \underline{H} build the 4-dimensional electromagnetic field force tensor $F_{ik} = (\underline{E}, \underline{H})$. The Maxwell stress tensor is given by

$$\sigma_{ik} = \frac{1}{4\pi} \left\{ -E_i E_k - H_i H_k + \frac{1}{2} \delta_{ik} (E^2 + H^2) \right\}.$$

With regards to "initial data for the Cauchy problem in general relativity" we refer to the corresponding lecture notes [PoD]:

Lecture 1: Introduction to Lorentzian geometry and causal theory

Lecture 2: The Einstein equations from a PDE perspective. The constraint equations and the local existence theorem of Choquet-Bruhat

Lecture 3: Solving the constraint equations via conformal method

Lecture 4: Topological censorship from the initial data point of view.

We further note the Weyl tensor representation

$$C_{iklm} = R_{iklm} - \frac{1}{2} R_{il} g_{km} + \frac{1}{2} R_{im} g_{kl} + \frac{1}{2} R_{kl} g_{im} - \frac{1}{2} R_{km} g_{il} + \frac{1}{6} R (g_{il} g_{km} - g_{im} g_{kl}) .$$

The reduced Einstein equations representation are given by ([PoD] lecture 2)

$$R_{\alpha,\alpha\beta}^H := -\frac{1}{2} g^{\gamma,\gamma} g_{\alpha\beta,\gamma\delta} + Q(g, \partial g) = 0 .$$

A "helicopter view" regarding Schrödinger's "purely quantum wave" vision

The Schrödinger (differentiation) operator is not bounded with respect to the norm of L_2 , i.e. only on a dense subspace of L_2 a corresponding spectral representation of this operator can be defined. The not vanishing constant Fourier term of the baseline Hermite polynomial (which is the Gaussian function) leads to mathematical challenges with respect to the creation and annihilation operators of the related Hamiltonian operator of the quantum oscillator model. The Hilbert transform of a function f has always vanishing constant Fourier terms. As a consequence, the Hilbert-transformed Schrödinger operator form with extended domain $H_{-1/2}$ is bounded (with respect to the norm of L_2) leading to a bounded Hermitian operator with corresponding spectral form representation.

Based on the newly defined common Hilbert space domain spectral theory can be applied, while

- the (physical) test space keeps the same, i.e. $L_2 = H_0$
- the current domains of the considered operators are extended to enable a (convergent) energy norm $\|x\|_{1/2}$ and a corresponding weak variation representation of the considered operator equations with respect to the inner product $(x, y)_{-1/2}$.

The corresponding notions from variation theory are "energy norm" and "operator norm" with correspondingly defined minimization problems ("energy" resp. "action" minimization problems). The corresponding eigenvalue problem of an operator T is then related to the inner product $(Tx, x)_{-1/2}$.

With respect to the newly proposed Pseudo-differential and Fourier multiplier operators with extended fractional Hilbert scale domain we note the following:

- The Maxwell equations are represented by differential equations or integral equations. Both representations are considered as equivalent.
- The Lagrange ("force") and the Hamiltonian ("energy") formalisms are considered as equivalent. The mathematical proof is based on the Legendre transform, i.e. the equivalence is only valid if the assumptions of the Legendre transform are fulfilled.

In both cases, corresponding (mathematical) regularity assumptions are required to enable those propositions. A restriction of the domain regularity of the considered operators leads to no longer well-defined classical differential equations resp. to no longer valid Lagrange formalism. In other words, the provided consistent model in the distributional framework represents the mathematical/transcendental view of the considered physical world, while the corresponding classical solutions of the several differential equations are mathematical approximations to those physical models. This concept also overcomes the "physical interpretation" challenge of the "Neumann PDE" representation of the pressure p in the NSE model.

As a consequence there is only a "one-energy" (field) concept and corresponding (PDE specific) manifestations/ forms of considered "Nature forces".

The proposed variation Hilbert space frame is built on the space-time frame with dimension $n = m + 1 = 4$. Therefore the **Huygens' Principle** (which is also valid for the initial value problem of the wave equation) is valid for all considered "wave" PDE, overcoming e.g. the $n > 10$ requirement of the string theory. At the same time, the characteristics roles of a space-time dimension = 4 is also underlined by the specific role of undistorted spherical travelling waves (Courant-Hilbert, "methods of mathematical physics", II, VI, §10.3).

Schrödinger's "purely quantum wave" vision is about half-odd integers, rather than integers to be applied to wave-mechanical vibrations which correspond to the motion of particles of a gas resp. the eigenvalues and eigen-functions of the harmonic quantum oscillator still governed by the Heisenberg uncertainty inequality. The alternatively proposed $H_{1/2}$ energy space enables Schrödinger's vision ([ScE] (7.23) ff):

let ω denotes the angular frequency, h the (h-bar) Planck constant and


$$e := \frac{\omega h}{2}$$

Then Schrödinger's "half-odd integer vision" is about the following replacement:

$n = 0$	$E_0 = e$	\rightarrow	$E_{1/2} = 1 * e$
$n > 0$	$E_1 = 1 * \omega h$	\rightarrow	$E_{3/2} = 2 * e$
	$E_2 = 2 * \omega h$	\rightarrow	$E_{5/2} = 3 * e$
\dots			
	$E_n = n * \omega h$	\rightarrow	$E_{(2n+1)/2} = (n + 1) * e, n = 0,1,2,\dots$

As a consequence the corresponding eigenvalue and eigenfunction solutions of the number operator (i.e. the product of generation and annihilation operators) start with index $n = 1$, not already with $n = 0$.

With respect to the ladder operators of the harmonic quantum Oscillator the proposed alternative quantum state and related energy Hilbert scales can be visualized by



Corresponding Hermite polynomials being linked to Weber's (Whittaker's) parabolic cylindrical polynomials by ([AbM] (13.1.32) [BuH] p. 215) are provided in section G.

We note that both, the Hermite polynomial system and the Hilbert transformed Hermite polynomial system build orthogonal systems of the Hilbert space $H_0 = L_2$. Therefore, generation and annihilation operators can be defined accordingly, whereby corresponding Hermite polynomials (original and Hilbert transform) are orthogonal with respect to the $H_0 = L_2$ inner product, as a consequence of the corresponding Hilbert transform property.

A "helicopter view" regarding a (truly infinitesimal geometry) Hilbert space based quantum gravity theory

The relativistic cosmology is based on the three assumptions, the cosmological principle, the Weyl postulate and the GRT. The GRT is based on the Riemann geometry with its underlying mathematical axiom, that the *Pythagoras theorem* is valid only for the case *when two points are infinitely near*.

We note that the Legendre (contact body) transform, which is applied to prove the equivalent of the Lagrange and Hamiltonian formalism, applied to differentials, is neglecting the $dxdy$ term in

$$d(x + y) = dx + dy \quad , \quad d(xy) = xdy + ydx + \mathbf{dxdy} = xdy + ydx$$

which is kind of contradiction to the Pythagoras axiom above.

The framework of the GRT is the affine connected manifolds ([WeH] p. 123). The metric character of a manifold is characterized relatively to a system of reference (= (1) *co-ordinate system* + (2) *calibration*) by two fundamental forms, namely a (1) quadratic differential form and a (2) linear one. They remain invariant during transformations to new co-ordinate systems.

The Weyl curvature tensor is a measure of a pseudo-Riemann manifold. It expresses the "tidal force" that a body feels when moving along a geodesic. It conveys the information how the shape of the body is distorted by this "tidal force". The Ricci curvature tensor, which expresses the trace components of the Riemann curvature tensor, contains the information about how the volumes change in the presence of "tidal forces", so the Weyl tensor is traceless component of the Riemann tensor. It is a tensor that has the same symmetries as the Riemann tensor with the extra condition that it be trace-free. The Weyl curvature is the only component of curvature for Ricci-flat manifolds and always governs the characteristics of the field equations of an Einstein manifold.

[WeH] p. 91: *The transition from Euclidean geometry to that of Riemann is founded in principle on the same idea as that which led from physics based on action at a distance to physics based on infinitely near action. The Ohm Law find by the observation, that the current flowing along a conducting wire is proportional to the difference of potential between the ends of the wire. Also the Coulomb Law deals with "actions at a distance". In order to model the physical model in its most general form, one accordingly deduces this law by reducing the measurements obtained to an infinitely small portion of wire. This results in the expression*

$$\text{curl}\vec{E} = 0 \quad , \quad \text{div}\vec{E} = \rho$$

on which Maxwell's theory is founded. Proceeding in the reverse direction, one derives from this differential law by mathematical processes the integral law, which we observed directly, on the supposition that conditions are everywhere similar (homogeneity). ... The fundamental fact of Euclidean geometry is that the square of the distance between two points is a quadratic form of the relative co-ordinates of the two points (Pythagoras Theorem).

We propose a replacement of the "Pythagoras theorem" in the infinitesimal small by the concept of "rotating differentials", leveraging on the Riesz operators property to be rotation invariant. This goes along with reduced regularity requirements to the corresponding operator domain. It also anticipate Leibniz' living force concept (see below).

At the same time it enables a replacement of the affine connected manifolds concept and its underlying invariant **exterior (covariant) derivative concept** (of p-forms) (e.g. enabling Hodge's potential theory of closed Riemann manifolds based on differential forms w/o vector fields) by a distributional Hilbert space concept with a corresponding **inner (differential) derivative product** (defining a corresponding (norm-) metric). The p-forms representations of the Riemann curvature tensor are e.g. given in ([FIH]). The Riemann geometry requires *differentiable* manifold w/o any physical meaning. The alternative (distributional) Hilbert space framework avoids this purely mathematical requirement, enabling also an alternative "orthogonality concept" as being applied in the Weyl postulate, where the world lines of the fluid particles, which act as the source of the gravitational field and which are often taken to model galaxies, should be *hypersurface orthogonal*.

In the context of the newly proposed "energy-space" $H_{1/2} = H_1 + H_1^-$ (where H_1^- represents the vacuum energy space) we note that already for the vacuum field equations ($R_{\alpha\beta} = 0$) there are two solutions, the Minkowski and the Schwarzschild metrics. Therefore those metrics are not compatible with the uniquely defined Hilbert space metric/norm. **Gödel's** example of a new type of cosmological solutions with non-vanishing density of matter (and with a cosmological term $\neq 0$) of Einstein's field equations provides a system with a rotation of matter relatively to the compass of inertia. This solution, or rather the properties of the four-dimensional space it defines are also provided ([GöK]).

The complementary variational analysis is proposed to characterize the solutions of the quantum gravitational field equations ([ArA]). The method of Noble is based on a system of two operator equations, which is analogue to the Euler differential equations, covering not only (non-linear) partial differential equations, but also integral equations. It is basically about a characterization of the solution as a saddle point of a minimization functional based on a "Hamiltonian" function $W(w, u)$, which is convex with respect to w , and concave with respect to u ([VeW] 6.2.4).

In continuum mechanics, the infinitesimal strain theory is about the deformation of a solid body. The displacement gradient is a 2nd order tensor, where it is possible to perform a geometric linearization of any one of the (infinitely many possible) strain tensors, e.g. the (Lagrange) strain tensor. Considering the linearized strain tensor as the "primary" unknown, instead of the displacement in the pure traction problem of three-dimensional linearized elasticity leads to a well-posed minimization problem, constrained by a weak form of the St Venant compatibility conditions. This approach also provides a new proof of Korn's inequality ([CiP]).

Regarding the NSE and YME the "infinitesimal small" of "fluids" and "quanta" is and will be all the time out of scope for any human observations. Mathematics is a purely descriptive science with well-established concepts to deal with any kind and "size" of "infinity" (e.g. Cauchy, Dedekind, Bolzano, Weierstrass, Kronecker, Cantor, Gödel, Brouwer). The mathematical tool managing physical "observations" are Partial Differential Equations (PDE), mathematical statistics and approximation theory. Those concepts are also applied in quantum mechanics and quantum field theory. The essential mathematical "objects" are the real numbers (while "nearly all" of those objects are far away from being "real") and the Lebesgue integral building the Lebesgue (Hilbert) space L_2 , where all rational numbers build a null set measured by its corresponding norm.

A "space lab" view comment regarding related philosophical concepts

We can think (hear and watch) the Yoda quote "may the FORCE be with us" and mathematics can model this FORCE/POWER/ENERGY in a way that all corresponding physical (law) models are consistent; ... the bad (or good?) news is, that's it and that's all! From a philosophical perspective we are back to

- Leibniz's ontology of force (e.g. "Fünf Schriften zur Logik und Metaphysik" and "Monadologie")

"the primitive active and passive forces, the form and matter are in the monadological view understood as features of the perceptions of the monads ... in this way the notion of force, ... loses its foundational status: primitive force gets folded into the perceptual life of non-extended perceiving things", Garber's monograph: Leibniz: Body, Substance, Monad, 2009)

- Kant's conception of physical matter and the existence of ether, which fills the whole space and time with its moving forces ([WoW])

"it is the moving forces of the ether that affect us"; ... "There exists a matter, distributed in the whole universe as a continuum, uniformly penetrating all bodies, and filling (all spaces) (thus not subject to displacement). Be it called ether, or caloric, or whatever...", ... "space is hypostatically", ... "Space which can be sensed (the object of the empirical intuition of space) is the complex of moving forces of matter - without which, space would be no object of possible experience", ... "Matter does not consist of simple parts, but each part is, in turn, composite...", Each part of matter is a quantum; i.e. matter does not consist of metaphysically simple parts, and Laplace's talk of material points (which were to be regarded as parts of matter) would, understood literally, contain a contradiction." ... "Atomism is a false doctrine of nature", ... forces "fill a space (both) extensively and intensively", ...

- Schrödinger's "(my) view of the world" with respect to "reasons for abandoning the dualism of thought and existence, or mind and matter"

"The objective world has only been constructed at the prize of taking the self, that is, mind, out of it remaking it; mind is not part of it; obviously, therefore, it can neither act on it nor be acted on by any of its parts. If this problem of the action of mind on matter cannot be solved within the framework of our scientific representation of the objective world, where and how can it be solved?" ... "No single man can make a distinction between the realm of his perceptions and the realm of things that cause it, since however detailed the knowledge he may have acquired about the whole world, the story is occurring only once and not twice. The duplication is an allegory suggested mainly by communication with other beings."

A TENTATIVE ANSWER: "A single experience that is never to repeat itself is biologically irrelevant. Biologic value lies only in learning the suitable reaction to a situation that offers itself again and again, in many cases periodically, and always requires the same response if the organism is to hold its ground." ... But whenever the situation exhibits a relevant differential - let us say the road is up at the place where we used to cross it, so that we have to make a detour - this differential and our response to it intrude into consciousness, from which, however, they soon fade below the threshold, if the differential becomes a constantly repeating feature. Now in those fashion differentials, variants of response, bifurcations, etc., are piled up one upon the other in unsurveyable abundance, but only the most recent ones remain in the domain of consciousness, only those with regard to which the living substance is still in the stage of learning or practicing.

... I would summarize my general hypothesis thus: consciousness is associated with the learning of living substance; it's knowing how (Können) is unconscious"

- Heidegger's notion of „mathematical“ physics ("Holzwege, die Zeit des Weltbildes" (72) ff: die neuzeitliche Physik heisst mathematische, weil sie ... eine ganz bestimmte Mathematik anwendet. ... Keineswegs wird das Wesen des Mathematischen durch das Zahlenhafte bestimmt. ... Durch sie (math. Physik) und für sie wird in einer betonten Weise etwas als das Schon-Bekannte im vorhinein ausgemacht. Der sich geschlossene Bewegungszusammenhang raum-zeitlich bezogener Massenpunkte. ... Kein Zeitpunkt hat vor einem anderen einen Vorzug. Jede Kraft bestimmt sich nach dem, ... was sie an Bewegung ..in der Zeiteinheit zur Folge hat.

A. The Riemann Hypothesis

All nontrivial zeros of the analytical continuation of the Riemann zeta function have a real part of 1/2. The Hilbert-Polya conjecture states that the imaginary parts of the zeros of the Zeta function corresponds to eigenvalues of an unbounded self adjoint operator.

We provide a solution for the RH building on a new Kummer function based Zeta function theory, alternatively to the current Gauss-Weierstrass function based Zeta function theory. This primarily enables a proof of the Hilbert-Polya conjecture (but also of other RH criteria like the Bagchi formulation of the Nyman-Beurling criterion or Polya criteria), whereby the imaginary parts of the zeros of the corresponding alternative Zeta function definition corresponds to eigenvalues of a bounded, self adjoint operator with (newly) distributional Hilbert space domain.

Let H and M denote the Hilbert and the Mellin transform operators. For the Gaussian function $f(x)$ it holds

$$M[f](s) = \frac{1}{2} \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) \quad , \quad M[-xf'(x)](s) = \frac{s}{2} \pi^{-s/2} \Gamma\left(\frac{s}{2}\right) = \frac{1}{2} \pi^{-s/2} \Pi\left(\frac{s}{2}\right)$$

The corresponding entire Zeta function is given by

$$\xi(s) := \frac{s}{2} \Gamma\left(\frac{s}{2}\right) (s-1) \pi^{-s/2} \zeta(s) = (1-s) \cdot \zeta(s) M[-xf'(x)](s) = \xi(1-s) \cdot$$

The central idea is to replace

$$M[-xf'(x)](s) \rightarrow M[f_H(x)](s)$$

by

$$M[f_H(x)](s) = 2\pi \cdot M\left[x {}_1F_1\left(1, \frac{3}{2}, -\pi x^2\right)\right](s) = \pi^{\frac{1-s}{2}} \Gamma\left(\frac{s}{2}\right) \tan\left(\frac{\pi}{2}s\right)$$

This enables the definition of an alternative entire Zeta function in the form

$$\xi^*(s) := (1-s) \pi^{\frac{1-s}{2}} \Gamma\left(\frac{s}{2}\right) \tan\left(\frac{\pi}{2}s\right) \cdot \zeta(s)$$

with same zeros as $\xi(s)$.

The fractional part function

$$\rho(x) := \{x\} := x - [x] = \frac{1}{2} - \sum_1^{\infty} \frac{\sin 2\pi vx}{\pi v} \in L_2^{\#}(0,1)$$

is linked to the Zeta function by

$$\zeta(1-s) = (s-1) M[\rho](s-1) = M[-x\rho'(x)](s-1)$$

The Hilbert transform of the fractional part function is given by

$$\rho_H(x) = \sum_1^{\infty} \frac{\cos 2\pi vx}{\pi v} = -\frac{1}{\pi} \log 2 \sin(\pi x) \in L_2^{\#}(0,1) \quad , \quad \hat{\rho}_H(0) = 0 \quad , \quad \rho_H' \in H_{-1}^{\#}$$

Applying the idea of above leads to the replacement

$$M[-x\rho'(x)](s-1) = \zeta(1-s) \quad \rightarrow \quad M[-\rho_H(x)](s-1)$$

$$M[-\rho_H(x)](s-1) = M\left[\sum_1^{\infty} \frac{\cos 2\pi vx}{\pi v}\right](s-1) = 2\pi \Gamma\left(\frac{1-s}{2}\right) \Gamma^{-1}\left(\frac{s}{2}\right) \cdot \chi(s) \cdot \frac{\zeta(s)}{s}$$

with same zeros as $\zeta(1-s)$.

The proposed framework also provides an answer to Derbyshire's question, ("Prime Obsession")

... "The non-trivial zeros of Riemann's zeta function arise from inquiries into the distribution of prime numbers. The eigenvalues of a random Hermitian matrix arise from inquiries into the behavior of systems of subatomic particles under the laws of quantum mechanics. What on earth does the distribution of prime numbers have to do with the behavior of subatomic particles?"

The answer, in a nutshell:

"identifying "fluids" or "sub-atomic particles" not with real numbers (scalar field, I. Newton), but with hyper-real numbers (G. W. Leibniz) enables a truly infinitesimal (geometric) distributional Hilbert space framework (H. Weyl) which corresponds to the Teichmüller theory, the Bounded Mean Oscillation (BMO) and the Harmonic Analysis theory. The distributional Hilbert scale framework enables the full power of spectral theory, while still keeping the standard $L(2)=H(0)$ -Hilbert space as test space to "measure" particles' locations. At the same time, the Ritz-Galerkin (energy or operator norm minimization) method and its counterpart, the methods of Trefftz/Noble to solve PDE by complementary variation principles (A. M. Arthurs, K. Friedrichs, L. B. Rall, P. D. Robinson, W. Velte) w/o anticipating boundary values) enables an alternative "quantization" method of PDE models (P. Ehrenfest), e.g. being applied to the Wheeler-de-Witt operator.

Regarding the proposed alternative quantization approach we also refer to the Berry-Keating conjecture. This is about an unknown quantization \mathbf{H} of the classical Hamiltonian $H=xp$, that the Riemann zeros coincide with the spectrum of the operator $1/2+i\mathbf{H}$. This is in contrast to canonical quantization, which leads to the Heisenberg uncertainty principle and the natural numbers as spectrum of the harmonic quantum oscillator. The Hamiltonian needs to be self-adjoint so that the quantization can be a realization of the Hilbert-Polya conjecture.

B. The Navier-Stokes Equations

The Navier-Stokes equations describe the motion of fluids. The *Navier–Stokes existence and smoothness* problem for the three-dimensional NSE, given some initial conditions, is to prove that smooth solutions always exist, or that if they do exist, they have bounded energy per unit mass.

We provide a global unique (weak, generalized Hopf) $H_{-1/2}$ -solution of the generalized 3D Navier-Stokes initial value problem. The global boundedness of a generalized energy inequality with respect to the energy Hilbert space $H_{1/2}$ is a consequence of the Sobolevskii estimate of the non-linear term (1959).

The "standard" weak Hopf solution is not well posed due to not appropriately defined domains of the underlying velocity and pressure operators. Therefore, this is also the case for the corresponding classical solution(s).

The proposed solution also overcomes the "Serin gap" issue, as a consequence of the bounded non-linear term with respect to the appropriate energy norm.

The Navier-Stokes Equations (NSE) describes a flow of incompressible, viscous fluid. The three key foundational questions of every PDE is existence, and uniqueness of solutions, as well as whether solutions corresponding to smooth initial data can develop singularities in finite time, and what these might mean. For the NSE satisfactory answers to those questions are available in two dimensions, i.e. 2D-NSE with smooth initial data possesses unique solutions which stay smooth forever. In three dimensions, those questions are still open. Only local existence and uniqueness results are known. Global existence of strong solutions has been proven only, when initial and external forces data are sufficiently smooth. Uniqueness and regularity of non-local Leray-Hopf solutions are still open problems.

Basically the existence of 3D solutions is proven only for "large" Banach spaces. The uniqueness is proven only in "small" Banach spaces. The question of global existence of smooth solutions vs. finite time blow up is one of the Clay Institute millennium problems.

The existence of weak solutions can be provided essentially by the energy inequality. If solutions would be classical ones, it is possible to prove their uniqueness. On the other side for existing weak solutions it is not clear that the derivatives appearing in the inequalities have any meaning.

Basically all existence proofs of weak solutions of the Navier-Stokes equations are given as limit (in the corresponding weak topology) of existing approximation solutions built on finite dimensional approximation spaces. The approximations are basically built by the Galerkin-Ritz method, whereby the approximation spaces are e.g. built on eigen-functions of the Stokes operator or generalized Fourier series approximations.

It has been questioned whether the NSE really describes general flows: The difficulty with ideal fluids, and the source of the d'Alembert paradox, is that in such fluids there are no frictional forces. Two neighboring portions of an ideal fluid can move at different velocities without rubbing on each other, provided they are separated by a streamline. It is clear that such a phenomenon can never occur in a real fluid, and the question is how frictional forces can be introduced into a model of a fluid.

The question intimately related to the uniqueness problem is the regularity of the solution. Do the solutions to the NSE blow-up in finite time? The solution is initially regular and unique, but at the instant T when it ceases to be unique (if such an instant exists), the regularity could also be lost. Given a smooth datum at time zero, will the solution of the NSE continue to be smooth and unique for all time?

There is no uniqueness proof for weak solutions except for over small time intervals. The simplest possible model example how a singularity can appear, is the ODE

$$y'(t) = y^2(t) \quad y(0) = y_0$$

with its solution

$$y(t) = \frac{y_0}{1 - t \cdot y_0}$$

which becomes infinite in finite time. For $n=3$ every positive solution of

$$y'(t) = cy^3(t)$$

blows up, i.e. there is no global estimate by this method.

The global boundedness of our solution is a consequence of the Sobolevskii-estimate of the non-linear term enabling the generalized energy inequality

$$\frac{1}{2} \frac{d}{dt} \|u\|_{-1/2}^2 + \|u\|_{1/2}^2 \leq |(Bu, u)_{-1/2}| \leq c \cdot \|u\|_{-1/2} \|u\|_1^2$$

Putting

$$y(t) := \|u\|_{-1/2}^2$$

one gets

$$y'(t) \leq c \cdot \|u\|_1^2 \cdot y^{1/2}(t)$$

This results into the a priori estimate

$$\|u(t)\|_{-1/2} \leq \|u(0)\|_{-1/2} + \int_0^t \|u\|_1^2(s) ds \leq c \left\{ \|u_0\|_{-1/2} + \|u_0\|_0^2 \right\}$$

ensuring global boundedness by the a priori energy estimate provided that $u_0 \in H_0$.

C. The Yang-Mills Equations

The YME are concerned with quantum field theory. Its related Millennium problem is about an appropriate mathematical model to govern the current "mass gap" of the YME, which is the difference in energy between the vacuum and the next lowest energy field.

The classical Yang-Mills theory is a generalization of the Maxwell theory of electromagnetism where the *chromo*-electromagnetic field itself carries charges. For given distributions of electric charges and currents the Maxwell equations determine the corresponding electromagnetic field. The laws by which the currents and charges behave are unknown. The energy tensor for electromagnetic fields is unknown for elementary particles. Matter is built by electromagnetic particles, but the field laws by which they are constituted are unknown, as well. The original inertia law (before Einstein's gravity theory) forced to attribute physical-objective properties to the space-time continuum. Analog to the Maxwell equations (in the framework of a short distance theory) Einstein considered the inertia law as a field property of the space-time continuum.

As a classical field theory the Maxwell equations have solutions which travel at the speed of light so that its quantum version should describe massless particles (gluons). However, the postulated phenomenon of color confinement permits only bound states of gluons, forming massive particles. This is the mass gap. Another aspect of confinement is asymptotic freedom which makes it conceivable that the quantum Yang-Mills theory exists without restriction to low energy scales. The problem is to establish rigorously the existence of the quantum Yang-Mills theory and a mass gap. The being challenged concept is about the concept of a *displacement current*, which is "just" about a mathematical requirement to enabled consistent mathematical data model.

Based on an $H_{1/2}$ energy Hilbert space we propose (analog to the NSE solution) a corresponding (weak) variation Maxwell equation representation. Its corresponding generalization (as described above) leads to a modified QED model. In the same manner as the Serrin gap issue has been resolved (as a result of the reduced regularity requirements) the chromo-electromagnetic field /particles can now carry charges. The open "field law" question above and how "particles" are interacting with each other to exchange energy are modeled in same manner as the coherent/incoherent turbulent flows of its NSE counterpart. The corresponding "zero state energy" model is no longer built on the Hermite polynomials but on its related Hilbert transformed Hermite polynomials, which also span the L_2 – Hilbert "test" space.

This provides a truly infinitesimal geometry (H. Weyl), enabling the concept of Riemann that force is a pseudo force only, which results from distortions of the geometrical structure. The baseline is a common Hilbert space framework (for all (nearby action) differential equations)

- providing the mathematical concept of a geometrical structure, while Riemann's manifold concept provides only a metric space and related affine connections
- building an integrated (no longer "force" dependent dynamical matter-field interaction laws) universal field model (including the gravity "force"), while replacing "force type" specific gauge fields and its combination model(s) for the electromagnetic, the strong and the weak nuclear power "forces"

As a consequence there are no "mass" and therefore no (YME-) "mass gap" anymore, but there is an appropriate vacuum (Hilbert) energy space, which is governed by the Heisenberg uncertainty principle.

The Heisenberg uncertainty inequality

Let $g \in H_0 = L_2(\mathbb{R})$, $\|g\|_0 = 1$, then

$$\mu(g) := t_0 \omega_0 := \left(\int_{\mathbb{R}} (t - t_0)^2 |g(t)|^2 dt \right) \left(\int_{\mathbb{R}} (\omega - \omega_0)^2 |\hat{g}(\omega)|^2 d\omega \right) \geq \frac{1}{4}$$

whereby t_0 denotes the mean value of the location of the particle and

$$\omega_0 := \int_{\mathbb{R}} \omega |\hat{g}(\omega)|^2 d\omega = \left(-i \frac{d}{dx} \varphi, \varphi \right)_0$$

the mean value of the momentum of the particle. The localization "uncertainty" $\mu(g)$ of the function g at the phase point (t_0, ω_0) is defined, provided that $-\infty < t_0, \omega_0 < \infty$, i.e. $g \in H_{1/2}$.

The proposed alternative quantum state Hilbert space $H_{-1/2}$ provides an alternative concept to the "Dirac function" calculus. This overcomes current handicaps concerning the regularity of the Dirac function, which depends from the space dimension, i.e. $\delta \in H_{-s}(\mathbb{R}^n)$ for $s > n/2$.

The alternatively proposed Hilbert space $H_{-1/2}$ provides a truly "microscopic" mathematical frame (independently from the space dimension), while still supporting the existing physical observation (statistical analysis) subspace. It is also proposed to replace the (continuous & differentiable) manifold concept (and exterior products of differential forms) in Einstein's field theory.

D. The Boltzmann Entropy

The Boltzmann equation is a (non-linear) integro-differential equation which forms the basis for the kinetic theory of gases. This not only covers classical gases, but also electron /neutron /photon transport in solids & plasmas / in nuclear reactors / in super-fluids and radiative transfer in planetary and stellar atmospheres. The Boltzmann equation is derived from the Liouville equation for a gas of rigid spheres, without the assumption of "molecular chaos"; the basic properties of the Boltzmann equation are then expounded and the idea of model equations introduced. Related equations are e.g. the Boltzmann equations for polyatomic gases, mixtures, neutrons, radiative transfer as well as the Fokker-Planck and Vlasov equations. The treatment of corresponding boundary conditions leads to the discussion of the phenomena of gas-surface interactions and the related role played by proof of the Boltzmann H-theorem.

We provide the relationships of the Boltzmann and Landau equations to the proposed alternative quantum state Hilbert space frame $H_{-1/2}$ resp. its corresponding quantum energy space $H_{1/2}$ and the related NSE global solution, as well as to the proposed alternative Schrödinger (Calderón) momentum operator. At the same point in time we suggest to apply the same frame for a proof of the non-linear Landau damping for the Vlasov equation with only physical relevant mathematical assumptions.

In line with the proposed distributional $H_{-1/2}$ -Hilbert space concept of this paper, we suggest to define "continuous entropy" in a weak $H_{-1/2}$ - frame in the form

$$h(X) := (f, \log \frac{1}{f})_{-1/2} ,$$

where X denotes a continuous random variable with density $f(x)$. In this case it can be derived from a Shannon (discrete) entropy in the limit of n , the number of symbols in distribution $P(x)$ of a discrete random variable X ([MaC]):

$$H(X) := \sum_i P(x_i) \log \left(\frac{1}{P(x_i)} \right) .$$

This distribution $P(x)$ can be derived from a set of axioms. This is not the case, in case of the standard entropy in the form

$$h(X) := (f, \log \frac{1}{f})_0 .$$

E. Wavelets

Wavelets are proposed as appropriate analysis tool for the proposed NMEP, additionally to Fourier analysis technique. There are at least two approaches to wavelet analysis, both are addressing the somehow contradiction by itself, that a function over the one-dimensional space \mathbb{R} can be unfolded into a function over the two-dimensional half-plane.

A wavelet transform W_ϑ

$$W_\vartheta[f](a, b) := |a|^{-1/2} \int_{\mathbb{R}} f(t) \vartheta\left(\frac{t-b}{a}\right) dt$$

of a function f with respect to a wavelet function ϑ is an isometric mapping. The admissibility condition is given by

$$0 < c_\vartheta := \int_{\mathbb{R}} \frac{|\hat{\vartheta}(\omega)|^2}{|\omega|} d\omega < \infty$$

The corresponding adjoint operator W_ϑ^* is given by the inverse wavelet transform on its range:

$$W_\vartheta^*[g](a, b) := c_\vartheta^{-1/2} \int_{\mathbb{R}} \int_{\mathbb{R}} |a|^{-1/2} g(a, b) \frac{1}{a} \vartheta\left(\frac{t-b}{a}\right) \frac{da}{a} db$$

The Fourier transform of a wavelet transformed function f is given by

$$\widehat{W_\vartheta[f]}(a, \omega) := (2\pi|a|)^{\frac{1}{2}} c_\vartheta^{-\frac{1}{2}} \hat{\vartheta}(-a\omega) \hat{f}(\omega) .$$

For $\varphi, \vartheta \in L_2(\mathbb{R})$, $f_1, f_2 \in L_2(\mathbb{R})$,

$$0 < |c_{\vartheta\varphi}| := 2\pi \left| \int_{\mathbb{R}} \frac{\hat{\vartheta}(\omega) \overline{\hat{\varphi}(\omega)}}{|\omega|} d\omega \right| < \infty$$

and $|c_{\vartheta\varphi}| \leq c_\vartheta c_\varphi$ one gets the duality relationship ([LoA])

$$(W_\vartheta f_1, W_\varphi^* f_2)_{L_2(\mathbb{R}^2, \frac{dadb}{a^2})} = c_{\vartheta\varphi} (f_1, f_2)_{L_2}$$

i.e.

$$W_\varphi^* W_\vartheta [f] = c_{\vartheta\varphi} f \quad \text{in a } L_2 \text{ -sense.}$$

This identity provides an additional degree of freedom to apply wavelet analysis with appropriately (problem specific) defined wavelets in a (distributional) Hilbert scale framework where the "microscope observations" of two wavelet (optics) functions ϑ, φ can be compared with each other by the above "reproducing" ("duality") formula. The prize to be paid is about additional efforts, when re-building the reconstruction wavelet. The extended admissibility condition above indicates that wavelet "pairs" in the form $(\varphi, \vartheta) \in L_2 \times H_{-1}$ would be an appropriate good baseline to start from, when analyzing in the Hilbert space frame $H_{-1/2} = L_2 \times L_2^-$, where L_2^- denotes the complementary space of L_2 with respect to the $H_{-1/2}$ -norm. The Hilbert transform operator (which is valid for every Hilbert scale) is the "natural" partner to the wavelet-transform operator, as it is skew-symmetric, rotation invariant and each Hilbert transformed "function" has vanishing constant Fourier term. The example in the context above is the Hilbert transform of the Gaussian distribution function, the (odd) Dawson function, with the "polynomial degree" point of zero at +/- infinite.

F. The (Hilbert) transform of a differential du (Plemelj) and the corresponding positive definite continued fraction (Stieltjes) transform representation of u

The Stieltjes continued fraction theory provides an integral representation of the continued fraction

$$\frac{1}{|a_1-z|} - \frac{b_1^2}{|a_2-z|} - \frac{b_2^2}{|a_3-z|} - \dots \quad , \quad a_n \in R, b_n \in R - \{0\}, z \in Z$$

in the form ([HeE], [BrK1])

$$u(z) = S[\sigma](z) := \int_{-\infty}^{\infty} \frac{d\sigma(\mu)}{z - \mu} = \frac{1}{|a_1-z|} - \frac{b_1^2}{|a_2-z|} - \frac{b_2^2}{|a_3-z|} - \dots$$

provided that the series $\sum_{n=1}^{\infty} |p_n(z)|^2$ diverge for at least one non-real $z \in Z - R$ (and therefore for all $z \in Z - R$) whereby the polynomials $p_n(z)$ are defined by the linear homogenous equations

$$(a_1 - z)p_1(z) - b_1 p_2(z) = 0$$

$$-b_{n-1} p_{n-1}(z) + (a_n - z)p_n(z) - b_n p_{n+1}(z) = 0 \quad .$$

The above is about a real bounded J-fraction ([WaH], theorem 27.4). The equivalent function of a positive definite J-fraction can be represented as a Stieltjes transform ([WaH], theorems 65.1 & 66.1).

Just as a side remark we note the result from P. Bundschuh: the number

$$\sum_{n=0}^{\infty} \frac{1}{n^2 + 1} = \frac{1}{2} + \frac{\pi e^{\pi} + e^{-\pi}}{2 e^{\pi} - e^{-\pi}} = \frac{1}{2} + \frac{\pi}{2} \coth(\pi)$$

is transcendental.

The above formula is related to the series ([ChK] VI, §2)

$$\frac{1}{2} - \frac{\pi}{2} z \cot(\pi z) = \frac{1}{2} + \frac{\pi}{2} z \left(\frac{d}{dz} \rho_H(z) \right) = \sum_{n=1}^{\infty} \zeta(2n) z^{2n} \quad , \quad |z| < 1 \quad .$$

G. Some further formulas

The half-odd integers are related to the Fourier coefficients of the convolution (integral) equation

$$Gu = f$$

where

$$G[u](y) := \int_{-\infty}^{\infty} g(y-x)u(x)dx, \quad g(x) := \frac{1}{\cosh(x)}$$

with the secans hyperbolicus function ([GrI], 1.232, 1.411)

$$\begin{aligned} g(x) := \operatorname{sech}(x) &:= \frac{1}{\cosh(x)} = \operatorname{cn}(x; 1) = \operatorname{dn}(x; 1) = 2 \sum_{n=0}^{\infty} (-1)^n e^{-(2n+1)x} \\ &= -2 \sum_{n=0}^{\infty} (-1)^{n-1} \frac{(n + \frac{1}{2})\pi}{(n + \frac{1}{2})^2\pi^2 + x^2} = 1 + 2 \sum_{n=1}^{\infty} \frac{E_{2n}}{(2n)!} x^{2n} \end{aligned}$$

defining a distribution function similar to the normal distribution function.

The Mellin, the Hilbert and the Fourier transforms of $g(x)$ are given by ([GrI], 3.523, 3.981)

$$M[g](s) = \int_0^{\infty} g(t)t^s \frac{dt}{t} = 2^{1-s}\gamma(s) {}_1F_1(-1; s; \frac{1}{2}), \quad (\text{where } \gamma \text{ denotes the Gamma function})$$

$$\begin{aligned} H[g](x) &= 2\pi \int_0^{\infty} \frac{\sin(2\pi xy)}{\coth(\frac{\pi}{2}y)} dy = \frac{1}{\pi} \frac{\sinh(4x)}{1+\cosh(4x)} = \frac{1}{\pi} \frac{\cosh(4x)-1}{\sinh(4x)} \\ &= -2 \sum_{n=0}^{\infty} (-1)^{n-1} \frac{x}{(n + \frac{1}{2})^2\pi^2 + x^2} \end{aligned}$$

$$\begin{aligned} F[g](x) := \hat{g}(\omega) &:= \int_{-\infty}^{\infty} g(t)e^{-i\omega t} dt = \int_{-\infty}^{\infty} \frac{e^{(1-i\omega)t}}{e^{2t}+1} dt = 2 \frac{\frac{\pi}{2}}{\cosh(\frac{\pi}{2}\omega)} \\ &= \pi g(\frac{\pi}{2}\omega) = -2 \sum_{n=0}^{\infty} (-1)^{n-1} \frac{(n + \frac{1}{2})}{(n + \frac{1}{2})^2 + (\frac{1}{2}\omega)^2} \end{aligned}$$

where

$$z_n := -i\pi(\frac{1}{2} + n), \quad n \in \mathbb{Z}$$

are the simple poles of the integrand above with the residuals

$$\operatorname{Res}_{z=z_n} \frac{e^{(1-i\omega)z}}{e^{2z}+1} = \frac{e^{(1-i\omega)z_n}}{e^{2z_n}+1} = \frac{i}{2} (-1)^n e^{-\pi\omega(n+\frac{1}{2})}$$

The solution of the above integral equation is then given by

$$u(x) = \frac{1}{4\pi} \left[\frac{2}{\pi} \cosh(\frac{\pi}{2}\omega) \hat{f}(\omega) \right] (-x).$$

The generalized Hermite polynomials satisfy different differential equations for even and odd polynomials enabling corresponding spectral analysis ([KrA]). For the special case of the Schrödinger differential equation the spectrum of its related Schrödinger equation operator L is discrete, consisting of the odd integers. The corresponding eigen-functions form a complete orthogonal set in the weighted- L_2 space ([DaD]).

The Galois group of the equations $K_n^{(i)}(x) = 0$, $i = 0,1$, is the symmetric group S_n , where

$$K_n^{(0)}(x^2) = H_{2n}(x) \quad , \quad xK_n^{(1)}(x^2) = H_{2n+1}(x)$$

and $H_n(x)$ denote the Hermite polynomials ([ScW]), i.e. the polynomials $K_n^{(i)}(x)$ are non-affine.

Our alternatively proposed Schrödinger (Calderón) equation operator differs from the standard operator L by its combination with the Hilbert-transform operator H based on extended domain, which is an unitary operator with corresponding spectral theorem and a representation

$$H = \cos(A) + i * \sin(A) = e^{iA}$$

whereby A denotes a Hermitian operator with a corresponding spectrum on the unit circle. This enables a spectral representation of the alternatively proposed Schrödinger equation operator, whereby the L_2 space is governed by a discrete spectrum, while the corresponding complementary space of L_2 is governed by a continuous spectrum, modelling the "ground state zero" "eigen-differential" / "wave packages".

From [GrI] 1.317, 1.411, we recall the following identities

$$\frac{\pi}{2} \tan\left(\frac{\pi}{2}x\right) = \frac{\pi}{2} \cdot \frac{1 - \cos(\pi x)}{\sin(\pi x)} = \frac{\pi}{2} \cdot \frac{\sin(\pi x)}{1 + \cos(\pi x)}$$

$$\frac{\pi}{2} \tan\left(\frac{\pi}{2}x\right) = \frac{1}{x} \cdot \sum_{k=1}^{\infty} \frac{2^{2k}-1}{(2k)!} |B_{2k}| \cdot (\pi x)^{2k} \quad , \quad x^2 < 1 \quad .$$

The expansion of the \tan – resp. the $x \cot(x)$ – term in series of simple fractions are given by ([GrI] 1.421)

$$\frac{\tan \frac{\pi x}{2}}{\frac{\pi x}{2}} = \frac{8}{\pi^2} \sum_{k=1}^{\infty} \frac{1}{(2k-1)^2 - x^2} \quad , \quad -\frac{\pi}{2} (1-x) \cot\left(\frac{\pi}{2}(1-x)\right) = -1 + 2 \sum_{k=1}^{\infty} \frac{(1-x)^2}{4k^2 - (1-x)^2} \quad .$$

With respect to the relationship of the Zeta function and the $\pi x \cot(\pi x)$ function we further note the following formulas ([BeB] 5, [TiE] 4.14, [AnG] entry 3, entry 44)

- i. $\sum_{n>x} \frac{1}{n^s} = \frac{1}{2\pi x} \int_{x-i\infty}^{x+i\infty} x^{-s} (-\pi z \cot(\pi z)) dz$
- ii. $z(s) = \sum_{n \leq x} \frac{1}{n^s} - \frac{x^{1-s}}{1-s} + O(x^{-\sigma}) \quad , \quad \sigma \geq \sigma_0 > 0 \quad , \quad |t| \leq \frac{2\pi x}{c} \quad , \quad c > 1$
- iii. $x \coth(x) = 1 + \frac{x^2}{3} - \frac{x^2}{5} + \frac{x^2}{7} \dots = 1 + \frac{x^2}{3} - \frac{x^2}{9} \left[\frac{x^5}{5} + \frac{4 \cdot 5 x^2}{2 \cdot 3} + \frac{2 \cdot 3 x^2}{4 \cdot 5} + \frac{6 \cdot 7 x^2}{4 \cdot 5} + \frac{4 \cdot 5 x^2}{6 \cdot 7} + \dots \right]$
- iv. $\left(\frac{\pi}{2} s\right) \cdot \coth\left(\frac{\pi}{2} s\right) = 1 + \frac{s^2}{1} - \frac{1^2(s^2+1^2)}{3} + \frac{2^2(s^2+2^2)}{5} - \frac{3^2(s^2+3^2)}{7} \dots \dots \quad s \in C.$

We note that the periodical continuation of the log-Gamma-function $\log\gamma(s)$ with domain $(0,1)$ is $\in H_{-1/2}^\#(0,1)$, also anticipating the odd sin-Fourier terms in the form $\frac{\log(2\pi n)}{2\pi n} + \frac{\gamma}{2\pi n}$ ([GrI] 6.443). The log-Gamma-function is linked to the log-sin-function by the formula

$$\log\gamma(x) + \log\gamma(1-x) = \log(2\pi) + \log\frac{1}{2\sin(\pi x)} = \log(2\pi) + \sum_{k=1}^{\infty} \frac{\cos(2\pi kx)}{k}$$

$$\frac{\pi}{2} \cot\left(\frac{\pi}{2}x\right) - \frac{\pi}{2} \cot\left(\frac{\pi}{2}(x-1)\right) = \frac{2\pi}{2\sin(\pi x)}.$$

We briefly sketch the relationship of Plemelj's alternative potential definition, the Hilbert transform and the Laplacian equation, which may be solved using simple layer potential:

For the fundamental solution of the Laplacian

$$\gamma(s-t) := -\frac{1}{\pi} \log|\theta(s) - \theta(t)|$$

the potential

$$v(s) = \int \gamma(s-t)u(t)dt$$

is replaced by

$$v(s) = \int \gamma(s-t)du(t).$$

By partial integration one gets

$$v(s) = - \int \partial_t \gamma(s-t)u(t) dt$$

For smooth boundaries one gets

$$\partial_t \gamma(s-t) = \frac{1}{2\pi} \cot \frac{s-t}{2} + g(s,t)$$

where $g(s,t)$ is correspondingly regular.

Putting $a_n := (2\pi)^{-1/2}(-2)^n n!$, ($n = 0,1,2 \dots$) the Hermite polynomials are linked to Weber's (Whittaker's) parabolic cylindrical polynomials by ([AbM] (13.1.32) [BuH] p. 215)

$$M_{\left(n+\frac{1}{2}\right)-\frac{1}{4};\frac{1}{4}}(z) := \sqrt{2} \frac{a_n}{(2n)!} z^{\frac{1}{4}} e^{-\frac{z}{2}} H e_{2n}(\sqrt{2z}) = \frac{n!}{\gamma\left(n+\frac{1}{2}\right)} z^{\frac{1}{2}z-\frac{1}{4}} e^{-\frac{z}{2}} L_n^{-\frac{1}{2}}(z) = z^{\frac{1}{4}} e^{-\frac{z}{2}} M\left(-n, \frac{1}{2}; z\right)$$

$$M_{\left(n+\frac{1}{2}\right)+\frac{1}{4};\frac{1}{4}}(z) := 2 \frac{a_n}{(2n+1)!} z^{\frac{1}{4}} e^{-\frac{z}{2}} H e_{2n+1}(\sqrt{2z}) = \frac{n!}{\gamma\left(n+\frac{3}{2}\right)} z^{\frac{1}{2}z+\frac{1}{4}} e^{-\frac{z}{2}} L_n^{\frac{1}{2}}(z) = z^{-\frac{1}{4}} e^{-\frac{z}{2}} \left[x M\left(-n, \frac{3}{2}; z\right) \right]$$

resp. ([AbM] (13.6) ([GrI] (9.231)

$$M\left(-n, \frac{1}{2}; \frac{z^2}{2}\right) = \frac{n!}{(2n)!} \left(-\frac{1}{2}\right)^{-n} H e_{2n}(z)$$

$$x M\left(-n, \frac{3}{2}; \frac{z^2}{2}\right) = \frac{n!}{(2n+1)!} \left(-\frac{1}{2}\right)^{-n} H e_{2n+1}(z)$$

where

$$M_{\left(n+\frac{1}{2}\right)+\mu;\mu}(z) = \frac{z^{\frac{1}{2}-\mu} e^{\frac{z}{2}}}{(2\mu+1)(2\mu+2)\dots(2\mu+n)} \frac{d^n}{dz^n} (z^{n+2\mu} e^{-z}).$$

The Mellin transform of $h_{\rho,\sigma}(z) := e^{-\frac{z}{2}} M_{\sigma,\rho}(z)$ is given by ([GrI] (7.621)

$$\int_0^{\infty} x^s h_{\rho,\sigma}(s) \frac{dx}{x} = \frac{\gamma(1+2\sigma)\gamma(\rho-s)\gamma\left(\frac{1}{2}+\sigma+s\right)}{\gamma\left(\frac{1}{2}+\sigma+\rho\right)\gamma\left(\frac{1}{2}+\sigma-s\right)}, \quad \operatorname{Re}\left(-\frac{1}{2}-\sigma\right) < \operatorname{Re}(s) < \operatorname{Re}(\rho).$$

References

- [AbM] Abramowitz M., Stegun I. A., Handbook of Mathematical Functions, Dover Publications, Inc., New York, 1965
- [AmS] Amini S., On Boundary Integral Operators for the Laplace and the Helmholtz Equations and Their Discretization, Engineering Analysis with Boundary Elements, Elsevier, 1999
- [AnG] Andrews G. E., Berndt B. C., Jacobsen L., Lamphere R. L., The Continued Fractions Found in the Unorganized Portions of Ramanujan's Notebooks, American Mathematical Society, Vol. 99, No 477, 1992
- [ArA] Arthurs A. M., Complementary Variational Principles, Clarendon Press, Oxford, 1970
- [AzA] Aziz A. K., Kellogg R. B., Finite Element Analysis of a Scattering Problem, Math. Comp. Vol. 37, No 156, (1981) 261-272
- [AzC] Arcozzi N., Xinwei L., Riesz Transforms on Spheres, Mathematical Research Letters 4, 401-412, 1997
- [BeB] Berndt B. C., Andrews G. E., Ramanujan's Notebooks Part I, Springer Verlag, New York, Berlin, Heidelberg, Tokyo, 1989
- [BeM] Berry M. V., Riemann's Zeta Function: A Model for Quantum Chaos? In: Seligman T. H., Nishioka H. (Eds), Quantum Chaos and Statistical Nuclear Physics, Lecture Notes in Physics, Vol. 263, Springer Verlag, Berlin, Heidelberg, (1986) pp. 1-17
- [BiJ] Bittencourt J. A., Fundamentals of Plasma Physics, Springer Verlag, New York, Berlin, Heidelberg, Hong Kong, London, Milan, Paris, Tokyo, 2004
- [BiD] Bleecker D., Gauge Theory and Variational Principles, Dover Publications, Inc., Mineola, New York, 1981
- [BrK] Braun K., Interior Error Estimates of the Ritz Method for Pseudo-Differential Equations, Jap. Journal of Applied Mathematics, 3, 1, 59-72, 1986
- [BrK1] Braun K., A new ground state energy model, <http://www.quantum-gravitation.de>
- [BrK2] Braun K., An alternative Schroedinger (Calderón) momentum operator enabling a quantum gravity model, <http://www.quantum-gravitation.de>
- [BrK3] Braun K., Unusual Hilbert or Hölder space frames for the elementary particles transport (Vlasov) equation, <http://www.navier-stokes-equations.com>
- [BuH] Buchholtz H., The confluent Hypergeometric Function: with Special Emphasis on its Applications, Springer Verlag, Berlin, Heidelberg, 1969
- [CaD] Cardon D., convolution Operators and Zeros of Entire Functions, Proc. Amer. Math. Soc., Vol.130, No. 6, pp. 1725-1734, 2002
- [ChK] Chandrasekharan K., Elliptic Functions, Springer Verlag, Berlin, 1985
- [ChW] Chen W., Analytical solution of transient scalar wave and diffusion problems of arbitrary dimensionality and geometry by RBF wavelet series, 2001, arXiv:cs/0110055,

- [ChF] Chouly F., Heuer N., A Nitsche-based domain decomposition method for hypersingular integral equations, *Numer. Math.* 121 (2012) 705-729
- [CiP] Ciarlet P., G., Ciarlet Jr. P., Another approach to linearized elasticity and Korn's inequality, *C. R. Acad. Sci. Paris, Ser. 1* 339 (2004) 307-312
- [CoD] Colton D., Kress R., *Integral Equation Methods in Scattering Theory*, SIAM, Philadelphia, 2013
- [CoR] Courant R., Hilbert D., *Methods of Mathematical Physics II*, New York, Interscience Publishers, Inc., 1962
- [DaD] Dai D.Q., Han B., Jia R.-Q., Galerkin analysis for the Schrödinger equation by wavelets, *Journal of Mathematical Physics*, Vol. 45, No. 3 (2004)
- [DuR] Duffin R. J., Hilbert Transforms in Yukawan Potential Theory, *Proc. Nat. Acad. Sci.*, 69, 12, 3677-3679, 1972
- [EdH] Edwards H. M., *Riemann's Zeta Function*, Dover Publications, Inc., Mineola, New York, 1974
- [EiA] Einstein A., *Grundzüge der Relativitätstheorie*, Fried. Vieweg & Sohn, Braunschweig / Wiesbaden, 1922
- [EiA1] Einstein A., *The Foundation of the General Theory of Relativity*, Princeton University Press, Princeton, 1997
- [EsG] Eskin G., I., *Boundary Value Problems for Elliptic Pseudodifferential Equations*, American Mathematical Society, Providence, Rhode Island, 1981
- [FIH] Flanders H., *Differential Forms with Applications to the Physical Science*, Dover Publications, Inc., New York, 1963
- [FIG] Fichera G., Existence theorems in elasticity-boundary value problems of elasticity with unilateral constraints, *Encyclopedia of Physics* (S. Flügge, Chief Editor), Vol. Via/2 : *Mechanics of Solids II* (C. Truesdell, Editor), pp. 347-424, Springer Verlag, Berlin, 1972
- [FrH] Friedrich H., Cauchy Problem for the Conformal Vacuum Field Equations in General Relativity, *Commun. Mathe. Phys.* 91, 445-472 (1983)
- [FöE] Förster E., *The Cambridge Edition of the works of Immanuel Kant: Opus posthumum*, Cambridge University Press, 1993
- [GeR] Geroch K., Kronheimer E. H., Penrose R., Ideal points in space-time, *Proc. Roy. Soc. London*, A347, 545-567, 1972
- [GöK] Gödel K., An Example of a New Type of Cosmological Solutions of Einstein's Field Equations of Gravitation, *Review of Modern Physics*, Vol. 21, No 3, 447-450, (1949)
- [GrI] Gradshteyn I. S., Ryzhik I. M., *Table of Integrals Series and Products*, Fourth Edition, Academic Press, New York, San Francisco, London, 1965
- [HeE] Hellinger E., Zur Stieltjesschen Kettenbruchtheorie, *Math. Ann.*, 86, 19-29 (1922)

- [KrA] Krall A. M., Spectral analysis for the Generalized Hermite Polynomials, Trans. Amer. Math. Soc., Vol. 344, No. 1 (1994) pp. 155-172
- [HeW] Heisenberg W., Physikalische Prinzipien der Quantentheorie, BI Hochschultaschenbuch Bd. 1, Mannheim, Wien, Zürich, 1991
- [HeK] Hensel K., Leopold Kronecker's Werke, reprinted (Alkal. Paper) Chelsea Publishing Company, New York, 1968
- [HiP] Higgs P. W., Spontaneous Symmetry Breakdown without Massless Bosons, Physical Review, Vol. 145, No 4, p. 1156-1162, 1966
- [KyP] Kythe P. K., Fundamental Solutions for Differential Operators and Applications, Birkhäuser Verlag, Boston, 1996
- [LeN] Lerner N., A Note on the Oseen Kernels, A. Bove et. Al. (eds.), Advances in Phase Space Analysis of Partial Differential Equations, DOI 10.1007/978-0-8161-9_10, Birkhäuser Verlag, Boston, 2009
- [LiP] Lions P. L., Compactness in Boltzmann's equation via Fourier integral operators and applications, J. Math. Kyoto Univ. 34, 2, 391-427, 1994
- [LiP1] Lions P. L., On Boltzmann and Landau equations, Phil. Trans. R. Soc. Lond. A, 346, 191-204, 1994
- [LoA] Lifanov I. K., Poltavskii L. N., Vainikko G. M., Hypersingular Integral Equations and Their Applications, Chapman & Hall/CRC, Boca Raton, London, New York, Washington, D. C. 2004
- [LoA] Louis A. K., Maass P., Rieder A., Wavelets, Theorie und Anwendungen, B. G. Teubner Stuttgart, 1998
- [MaC] Marsh C., Introduction to Continuous Entropy, Department of Computer Science, Princeton University, December 13, 2013
- [MeY] Meyer Y., Coifman R., Wavelets, Calderón-Zygmund and multilinear operators, Cambridge studies in advanced mathematics 48, Cambridge University Press, 1996
- [MoC] Mouhot C., Villani C., On Landau damping, Acta Mathematica, Vol. 207, Issue 1, p. 29-201, 2011
- [PeB] Petersen B. E., Introduction to the Fourier Transform & Pseudo-differential Operators, Piman Advanced Publishing Program, Boston, London, Melbourne, 1983
- [PeR] Penrose R., Cycles of Time – An Extraordinary New View of the Universe, The Bodley Head, 2010
- [PIJ] Plemelj J. Potentialtheoretische Untersuchungen, B. G. Teubner Verlag, Leipzig, 1911
- [PoD] Pollack D., Initial Data for the Cauchy Problem in General Relativity, General Relativity Spring School 2015, Junior Scientist Andrejewski Days, March 22nd to April 4th, 2015, Brandenburg an der Havel, Germany

- [RuC] Runge C., Über eine Analogie der Cauchy-Riemannschen Differentialgleichungen in drei Dimensionen, Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Vol.1922, p. 129-136, 1922
- [ScE] Schrödinger E., Statistical Thermodynamics, Dover Publications, Inc., New York, 1989
- [ScW] Schulz W. Ueber die Galoissche Gruppe der Hermiteschen Polynome, J. reine und angew. Math., 177, 4, 248-252 (1937)
- [ShI] Shestakov I. V., Shlapunov A., On the Cauchy Problem for Operators with Injective Symbols in Sobolev Spaces
- [StA] Stern A., Tong Y., Desbrun M., Marsden J. E., Variational Integrators for Maxwell's Equations with Sources, PIERS ONLINE, Vol. 4, No. 7, 2008
- [StE] Stein E. M., Singular Integrals and Differentiability Properties of Functions, Princeton University Press, Princeton, New Jersey, 1970
- [TaN] Tarkhanov N. N., The Cauchy Problem for Solutions of Elliptic Equations, Akademie Verlag GmbH, Berlin 1995
- [TiE] Titchmarsh E. C., The Theory of the Riemann Zeta-function, Oxford University Press Inc., New York, first published 1951, Second Edition 1986
- [ViC] Villani C., Landau damping, Proceedings Of the International Congress Mathematicians, Hyderabad, India 2010
- [VeW] Welte W., Direkte Methoden der Variationsrechnung, Teubner Studienbücher, B. G. Teubner, Stuttgart, 1976
- [WaH] Wall H. S., Analytic Theory of Continued Fractions, AMS Chelsea Publishing, Providence, Rhode Island, reprinted by AMS, 2000
- [WeH] Weyl H., Space, Time, Matter, Dover Publications Inc., 1950
- [WeP] Werner P., Self-Adjoint Extensions of the Laplace Operator with Respect to Electric and Magnetic Boundary Conditions, J. Math. Anal. 70 (1979) 131-160
- [WeP1] Werner P., Spectral Properties of the Laplace Operator with respect to Electric and Magnetic Boundary Conditions, J. Math. Anal. 92 (1983) 1-65
- [WhJ] Whittaker J. M., Interpolatory Function Theory, Cambridge University Press, Cambridge, 1935
- [WoW] Wong W.-C., Kant's Conception of Ether as a Field in the Opus posthumum, in Ralph Schumacher, Rolf-Peter Horstmann & Volker Gerhardt (eds.), Kant Und Die Berliner Aufklärung, De Gruyter, pp. 676-684 (2001)