



Function spaces between BMO and critical Sobolev spaces[☆]

Jean Van Schaftingen

*Département de Mathématique, Université Catholique de Louvain, 2 chemin du Cyclotron,
1348 Louvain-la-Neuve, Belgium*

Received 29 November 2005; accepted 1 March 2006

Available online 2 May 2006

Communicated by H. Brezis

Abstract

The function spaces $D_k(\mathbf{R}^n)$ are introduced and studied. The definition of these spaces is based on a regularity property for the critical Sobolev spaces $W^{s,p}(\mathbf{R}^n)$, where $sp = n$, obtained by J. Bourgain, H. Brezis, New estimates for the Laplacian, the div-curl, and related Hodge systems, C. R. Math. Acad. Sci. Paris 338 (7) (2004) 539–543 (see also J. Van Schaftingen, Estimates for L^1 -vector fields, C. R. Math. Acad. Sci. Paris 339 (3) (2004) 181–186). The spaces $D_k(\mathbf{R}^n)$ contain all the critical Sobolev spaces. They are embedded in $BMO(\mathbf{R}^n)$, but not in $VMO(\mathbf{R}^n)$. Moreover, they have some extension and trace properties that $BMO(\mathbf{R}^n)$ does not have.

© 2006 Elsevier Inc. All rights reserved.

Keywords: Critical Sobolev spaces; BMO

1. Introduction

1.1. Integrals with divergence-free vector-fields

When $p < n$, the Sobolev space $W^{1,p}(\mathbf{R}^n)$ of functions whose distributional derivative is in $L^p(\mathbf{R}^n)$ is continuously embedded in the space $L^{p^*}(\mathbf{R}^n)$, with $p^* = np/(n - p)$, while when

[☆] The author is supported by a Research Fellow Grant of the Fonds National de la Recherche Scientifique. This work is partially supported by an EC Grant through the RTN Program “Front-Singularities” HPRN-CT-2002-00274.

E-mail address: vanschaftingen@math.ucl.ac.be.

$p > n$ it is embedded in the space of Hölder continuous functions of exponent α , $C^{0,\alpha}(\mathbf{R}^n)$, with $\alpha = 1 - n/p$ [1,5,19].

The case $p = n$ is more delicate. When $n > 1$, functions in $W^{1,n}(\mathbf{R}^n)$ do not need to be continuous or bounded, but have many properties in common with such functions. This is expressed for example by the embedding of $W^{1,n}(\mathbf{R}^n)$ in the spaces $BMO(\mathbf{R}^n)$ and $VMO(\mathbf{R}^n)$ of functions of bounded and vanishing mean oscillation [6]. These considerations are also valid for fractional Sobolev spaces $W^{s,p}(\mathbf{R}^n)$, with $sp = n$.

Another property of critical Sobolev space was recently obtained by Bourgain and Brezis [3, 23]: for every vector field $\varphi \in (L^1 \cap C)(\mathbf{R}^n; \mathbf{R}^n)$ and $u \in W^{s,p}(\mathbf{R}^n)$, if $\text{div } \varphi = 0$ in the sense of distributions, then

$$\left| \int_{\mathbf{R}^n} u \varphi \, dx \right| \leq C_{s,p} \|\varphi\|_{L^1(\mathbf{R}^n)} \|u\|_{W^{s,p}(\mathbf{R}^n)}. \tag{1.1}$$

There is no such property for $BMO(\mathbf{R}^n)$ or for $VMO(\mathbf{R}^n)$ (see [2] and Remark 5.2).

A natural question is the relationship between (1.1) and the embedding of $W^{s,p}(\mathbf{R}^n)$ in the spaces $BMO(\mathbf{R}^n)$ and $VMO(\mathbf{R}^n)$. In order to answer it, we define, for $n \geq 1$, the seminorm

$$\|u\|_{D_{n-1}(\mathbf{R}^n)} = \sup_{\substack{\varphi \in \mathcal{D}(\mathbf{R}^n; \mathbf{R}^n) \\ \text{div } \varphi = 0 \\ \|\varphi\|_{L^1(\mathbf{R}^n)} \leq 1}} \left| \int_{\mathbf{R}^n} u \varphi \, dx \right| \tag{1.2}$$

and the vector space

$$D_{n-1}(\mathbf{R}^n) = \{u \in \mathcal{D}'(\mathbf{R}^n) : \|u\|_{D_{n-1}(\mathbf{R}^n)} < \infty\}.$$

Here $\mathcal{D}(\mathbf{R}^N; \mathbf{R}^N)$ is the space of compactly supported smooth vector fields and $\mathcal{D}'(\mathbf{R}^n)$ is the space of distributions [16]. The subscript $n - 1$ will be justified by further extensions. By the inequality (1.1), $W^{s,p}(\mathbf{R}^n)$ is embedded in $D_{n-1}(\mathbf{R}^n)$.

The question of the previous paragraph is answered as follows: $VMO(\mathbf{R}^n)$ is not embedded in $D_{n-1}(\mathbf{R}^n)$ (Proposition 5.1), and $D_{n-1}(\mathbf{R}^n)$ is embedded in $BMO(\mathbf{R}^n)$ (Theorem 5.3). Moreover, if $u \in D_{n-1}(\mathbf{R}^n)$ is continuous, and $k \geq 2$, then $\|u\|_{\mathbf{R}^k} \|_{BMO(\mathbf{R}^k)} \leq C \|u\|_{D_{n-1}(\mathbf{R}^n)}$ (Theorems 3.4 and 5.3). This inequality remains open when $k = 1$.

The proof of the embedding of $D_{n-1}(\mathbf{R}^n)$ in $BMO(\mathbf{R}^n)$ is based on the duality between $BMO(\mathbf{R}^n)$ and the Hardy space $H^1(\mathbf{R}^n)$, and on a decomposition of every function in $H^1(\mathbf{R}^n)$ as a sum of some components of divergence-free vector-fields, with a suitable control on the norms.

The inequality (1.1) was preceded by a geometric counterpart [4]: for every closed rectifiable curve $\gamma \in C^1(S^1; \mathbf{R}^n)$ and $u \in (C \cap W^{1,n})(\mathbf{R}^n)$,

$$\left| \int_{\mathbf{R}^n} u(\gamma(t)) \dot{\gamma}(t) \, dt \right| \leq C_{s,p} \|\dot{\gamma}\|_{L^1(S^1)} \|u\|_{W^{s,p}(\mathbf{R}^n)}. \tag{1.3}$$

(See [22] for an elementary proof.) The right-hand side of (1.3) could also be used to define a seminorm on continuous functions. By the arguments of [3], based on a decomposition of divergence-free vector-fields in solenoids of Smirnov [18], one has in fact

$$\|u\|_{D_{n-1}(\mathbf{R}^n)} = \sup_{\gamma \in C^1(S^1; \mathbf{R}^n)} \frac{1}{\|\dot{\gamma}\|_{L^1(S^1)}} \left| \int_{S^1} u(\gamma(t)) \dot{\gamma}(t) dt \right|.$$

An open problem is whether restricting the curves on the right-hand side to be contained in k -dimensional planes, to triangles or to circles would yield an equivalent norm. The restriction to curves contained in k -dimensional planes is equivalent to requiring φ in (1.2) to have a range whose dimensions is at most k , see Section 6.4.

If $s \leq 1$, $sp = n$, and $u \in W^{s,p}(\mathbf{R}^n)$, then $u_+ \in W^{s,p}(\mathbf{R}^n)$. This property also holds in $BMO(\mathbf{R}^n)$. We do not know whether it holds for $D_{n-1}(\mathbf{R}^n)$. The question whether, for a given $\varphi : \mathbf{R} \rightarrow \mathbf{R}$ one has $\varphi(u) \in D_{n-1}(\mathbf{R}^n)$ whenever $u \in D_{n-1}(\mathbf{R}^n)$ remains open when φ is not affine.

1.2. Integrals with curl-free vector-fields

When $s = 1$ and $p = n = 2$, the inequality (1.1) is in fact a dual statement of the Sobolev–Nirenberg embedding

$$\|g\|_{L^{n/(n-1)}(\mathbf{R}^n)} \leq C \|Dg\|_{L^1(\mathbf{R}^n)}. \tag{1.4}$$

When $s = 1$ and $p = n > 2$, the estimate (1.1) is stronger than the embedding (1.4). If $n = 3$, (1.4) yields by duality that, for every $\varphi \in \mathcal{D}(\mathbf{R}^3; \mathbf{R}^3)$ and $u \in W^{1,3}(\mathbf{R}^3)$, if $\text{curl } \varphi = 0$ in the sense of distributions,

$$\left| \int_{\mathbf{R}^3} u \varphi dx \right| \leq C \|\varphi\|_{L^1(\mathbf{R}^3)} \|u\|_{W^n(\mathbf{R}^3)}. \tag{1.5}$$

For $u \in W^{s,p}(\mathbf{R}^3)$ with $sp = 3$, this inequality can be deduced from (1.1) recalling that, for every $e \in \mathbf{R}^3$

$$\text{div}(\varphi \times e) = (\text{curl } \varphi) \cdot e. \tag{1.6}$$

In \mathbf{R}^3 , one can now investigate the relationship between (1.1), (1.5), and the embedding of $W^{s,p}(\mathbf{R}^n)$ in the spaces $BMO(\mathbf{R}^n)$ and $VMO(\mathbf{R}^n)$. We define therefore the seminorm

$$\|u\|_{D_1(\mathbf{R}^3)} = \sup_{\substack{\varphi \in \mathcal{D}(\mathbf{R}^3; \mathbf{R}^3) \\ \text{curl } \varphi = 0 \\ \|\varphi\|_{L^1(\mathbf{R}^3)} \leq 1}} \left| \int_{\mathbf{R}^3} u \varphi dx \right|$$

and the vector space

$$D_1(\mathbf{R}^3) = \{u \in \mathcal{D}'(\mathbf{R}^3) : \|u\|_{D_1(\mathbf{R}^3)} < \infty\}.$$

While $VMO(\mathbf{R}^3) \not\subset D_1(\mathbf{R}^3)$, one has the following continuous embeddings:

$$D_2(\mathbf{R}^3) \subset D_1(\mathbf{R}^3) \subset BMO(\mathbf{R}^3).$$

The first embedding is a consequence of (1.6), and the second of the duality between $BMO(\mathbf{R}^3)$ and the Hardy space $H^1(\mathbf{R}^3)$, and of a decomposition of every function in $H^1(\mathbf{R}^3)$ as a sum of some components of curl-free vector-fields.

If $u \in D_1(\mathbf{R}^2)$, its extension $U(x, y) = u(x)$ to \mathbf{R}^3 is in $D_1(\mathbf{R}^3)$. It would be in $D_1(\mathbf{R}^3)$ if and only if U was bounded. On the other hand, if $u \in D_2(\mathbf{R}^3)$ is continuous, one has the trace inequality $\|u|_{\mathbf{R}^2}\|_{D_1(\mathbf{R}^2)} \leq C \|u\|_{D_2(\mathbf{R}^3)}$. The problem whether the trace inequalities $\|u|_{\mathbf{R}^2}\|_{BMO(\mathbf{R}^2)} \leq C \|u\|_{D_1(\mathbf{R}^3)}$ and $\|u|_{\mathbf{R}}\|_{BMO(\mathbf{R})} \leq C \|u\|_{D_2(\mathbf{R}^3)}$ hold is open.

The seminorm $\|\cdot\|_{D_1(\mathbf{R}^3)}$ can also be characterized geometrically: by the co-area formula, for every $u \in C(\mathbf{R}^3)$,

$$\|u\|_{D_1(\mathbf{R}^3)} = \sup_{\Omega} \frac{1}{\mathcal{H}^2(\partial\Omega)} \left| \int_{\partial\Omega} u(y)v(y) d\mathcal{H}^2(y) \right|,$$

where the supremum is taken over bounded domains $\Omega \subset \mathbf{R}^3$ with a smooth connected boundary, $v(y)$ is the unit exterior normal vector to the boundary at $y \in \partial\Omega$, and \mathcal{H}^2 is the two-dimensional Hausdorff measure.

1.3. Integrals along differential forms

In higher dimensions, the generalization of (1.1) corresponding to (1.5) in \mathbf{R}^3 is expressed with differential forms: if $1 \leq k \leq n - 1$, then, for every compactly supported smooth k -differential form $\varphi \in \mathcal{D}(\mathbf{R}^n; \Lambda^k \mathbf{R}^n)$ and for every $u \in W^{s,p}(\mathbf{R}^n)$ with $p \geq 1$ and $sp = n$, if $d\varphi = 0$, then

$$\left| \int_{\mathbf{R}^n} u\varphi dx \right| \leq C_{s,p} \|\varphi\|_{L^1(\mathbf{R}^n)} \|u\|_{W^{s,p}(\mathbf{R}^n)}. \tag{1.7}$$

The previous definitions of $D_k(\mathbf{R}^n)$ are generalized as follows. For $1 \leq k \leq n - 1$, we define the seminorm

$$\|u\|_{D_k(\mathbf{R}^n)} = \sup_{\substack{\varphi \in \mathcal{D}(\mathbf{R}^n; \Lambda^k \mathbf{R}^n) \\ d\varphi=0 \\ \|\varphi\|_{L^1(\mathbf{R}^n)} \leq 1}} \left| \int_{\mathbf{R}^n} u\varphi dx \right|$$

and the vector space

$$D_k(\mathbf{R}^n) = \{u \in \mathcal{D}'(\mathbf{R}^n): \|u\|_{D_k(\mathbf{R}^n)} < \infty\}.$$

By (1.7), $W^{s,p}(\mathbf{R}^n) \subset D_k(\mathbf{R}^n)$. These spaces $D_k(\mathbf{R}^n)$ also contain other functions, such as $\log(\sum_{i=1}^{k+1} x_i^2)$.

The spaces $D_k(\mathbf{R}^n)$ contain neither $BMO(\mathbf{R}^n)$ nor $VMO(\mathbf{R}^n)$. Our main result is that $D_k(\mathbf{R}^n)$ is embedded in $BMO(\mathbf{R}^n)$. We first show that $D_k(\mathbf{R}^n)$ is embedded in $D_1(\mathbf{R}^n)$, then we prove

that $D_1(\mathbf{R}^n)$ is embedded in $BMO(\mathbf{R}^n)$, in the same fashion as the embedding of $D_1(\mathbf{R}^3)$ in $BMO(\mathbf{R}^3)$ outlined above.

The other known properties of the spaces $D_k(\mathbf{R}^n)$ can be summarized as follows. The seminorm $\|\cdot\|_{D_k(\mathbf{R}^n)}$ is a norm modulo constants and the space $D_k(\mathbf{R}^n)$ modulo constants is a Banach space. The spaces $D_k(\mathbf{R}^n)$ are all different and are decreasing with respect to k . The spaces $D_k(\mathbf{R}^n)$ also have a trace property. If $u \in D_K(\mathbf{R}^N)$ is a limit of continuous functions, then it has a well-defined trace in $D_k(\mathbf{R}^n)$ if $N - K = n - k$. On the other hand, the function spaces $D_k(\mathbf{R}^n)$ do not have better integrability properties than the exponential integrability of functions in $BMO(\mathbf{R}^n)$.

1.4. Organization of the paper

We define in Section 2 the spaces $D_k(\mathbf{R}^n)$ for $1 \leq k \leq n$ by duality on closed smooth forms, and characterize them by duality on exact forms (Proposition 2.6). The space $D_n(\mathbf{R}^n)$ is in fact $L^\infty(\mathbf{R}^n)/\mathbf{R}$ (Proposition 2.9). We characterize geometrically the seminorm for continuous functions in the cases $k = 1$ (Proposition 2.10) and $k = n - 1$ (Proposition 2.11). For $1 < k < n - 1$, there is an equivalent seminorm defined by integration on real polyhedral chains without boundary (Proposition 2.14).

Section 3 reviews the basic properties of the spaces $D_k(\mathbf{R}^n)$: mutual injections (Theorem 3.1), extension theory (Theorem 3.2) and trace theory (Theorem 3.4).

Section 4 gives examples of functions in $D_k(\mathbf{R}^n)$: critical Sobolev spaces (Theorem 4.1) and other functions (Propositions 4.3, 4.6 and 4.10).

The embedding of the spaces $D_k(\mathbf{R}^n)$ in $BMO(\mathbf{R}^n)$ is proved in Section 5 (Theorem 5.3). This provides an easy proof of the completeness of $D_k(\mathbf{R}^n)$ (Theorem 5.5). We also address the question of improved integrability of functions in $D_k(\mathbf{R}^n)$ (Proposition 5.6).

The paper ends with considerations about further problems in the study of the spaces $D_k(\mathbf{R}^n)$ (Section 6).

Appendix A is devoted to density properties of closed and exact smooth forms.

2. Definitions and characterizations

2.1. Preliminaries

The space of k -forms on \mathbf{R}^n is denoted by $\Lambda^k \mathbf{R}^n$. The exterior product of $\alpha \in \Lambda^k \mathbf{R}^n$ and $\beta \in \Lambda^\ell \mathbf{R}^n$ is $\alpha \wedge \beta \in \Lambda^{k+\ell} \mathbf{R}^n$. The space $\Lambda^1 \mathbf{R}^n$ is the dual of \mathbf{R}^n and has a canonical basis $\omega_1, \dots, \omega_n$ biorthogonal to the canonical basis e_1, \dots, e_n of \mathbf{R}^n . Moreover, $\Lambda^k \mathbf{R}^n$ has a canonical Euclidean norm denoted by $|\cdot|$.

A differential form is a function $\varphi: \mathbf{R}^n \rightarrow \Lambda^k \mathbf{R}^n$. The exterior differential d is defined by

$$d\varphi = \sum_{i=1}^n \omega_i \wedge \frac{\partial \varphi}{\partial x_i}.$$

(This makes sense if e.g. φ is a C^1 function or a distribution.)

If V is finite-dimensional, the space of V -valued compactly supported smooth (C^∞) functions (test functions) is denoted by $\mathcal{D}(\mathbf{R}^n; V)$ and is endowed with its usual topology [16]. The space of distributions $\mathcal{D}'(\mathbf{R}^n)$ is the dual of $\mathcal{D}(\mathbf{R}^n) = \mathcal{D}(\mathbf{R}^n; \mathbf{R})$.

Lebesgue’s measure on \mathbf{R}^n is denoted by \mathcal{L}^n and the r -dimensional Hausdorff measure by \mathcal{H}^r .

2.2. Definition

The spaces $D_k(\mathbf{R}^n)$ are defined in terms of appropriate test function spaces.

Definition 2.1. For $1 \leq k \leq n$, define

$$D_{\#}(\mathbf{R}^n; \Lambda^k \mathbf{R}^n) = \left\{ \varphi \in \mathcal{D}(\mathbf{R}^n; \Lambda^k \mathbf{R}^n) : d\varphi = 0 \text{ and } \int_{\mathbf{R}^n} \varphi \, dx = 0 \right\},$$

$$L_{\#}^1(\mathbf{R}^n; \Lambda^k \mathbf{R}^n) = \left\{ \varphi \in L^1(\mathbf{R}^n; \Lambda^k \mathbf{R}^n) : d\varphi = 0 \text{ and } \int_{\mathbf{R}^n} \varphi \, dx = 0 \right\}.$$

Remark 2.2. The restriction $1 \leq k \leq n$, is justified by the fact that

$$L_{\#}^1(\mathbf{R}^n; \Lambda^k \mathbf{R}^n) = D_{\#}(\mathbf{R}^n; \Lambda^k \mathbf{R}^n) = \{0\}$$

when $k = 0$ or $k > n$.

Remark 2.3. If $1 \leq k \leq n - 1$, $\varphi \in L^1(\mathbf{R}^n; \Lambda^k \mathbf{R}^n)$ and $d\varphi = 0$, then $\int_{\mathbf{R}^n} \varphi = 0$, while for every $\varphi \in L^1(\mathbf{R}^n; \Lambda^n \mathbf{R}^n)$, $d\varphi = 0$. Therefore, for a given $1 \leq k \leq n$, only one condition in the definition is essential.

Definition 2.4. For $1 \leq k \leq n$, and $u \in \mathcal{D}'(\mathbf{R}^n)$, let

$$\|u\|_{D_k(\mathbf{R}^n)} = \sup_{\substack{\varphi \in D_{\#}(\mathbf{R}^n; \Lambda^k \mathbf{R}^n) \\ \|\varphi\|_{L^1(\mathbf{R}^n)} \leq 1}} \left| \int_{\mathbf{R}^n} u \varphi \, dx \right|,$$

and define

$$D_k(\mathbf{R}^n) = \{u \in \mathcal{D}'(\mathbf{R}^n) : \|u\|_{D_k(\mathbf{R}^n)} < \infty\}.$$

The integral appearing in the definition of $\|\cdot\|_{D_k(\mathbf{R}^n)}$ should be understood as a duality product between $\mathcal{D}'(\mathbf{R}^n)$ and $\mathcal{D}(\mathbf{R}^n; \Lambda^k \mathbf{R}^n)$, which takes its values in $\Lambda^k \mathbf{R}^n$, while $|\cdot|$ is the standard Euclidean norm of this integral. The set $D_k(\mathbf{R}^n)$ is a vector space; the function $\|\cdot\|_{D_k(\mathbf{R}^n)}$ is a seminorm on $D_k(\mathbf{R}^n)$, and vanishes for constant distributions.

Remark 2.5. In fact, by Theorem 5.3, if $\|u\|_{D_k(\mathbf{R}^n)} = 0$, then $\|u\|_{BMO(\mathbf{R}^n)} = 0$, whence u is constant.

The seminorm $\|\cdot\|_{D_k(\mathbf{R}^n)}$ can also be computed by considering exterior differentials of compactly supported smooth forms in place of closed compactly supported smooth forms.

Proposition 2.6. Let $1 \leq k \leq n$. For every $u \in \mathcal{D}'(\mathbf{R}^n)$,

$$\|u\|_{D_k(\mathbf{R}^n)} = \sup_{\substack{\psi \in \mathcal{D}(\mathbf{R}^n; \Lambda^{k-1} \mathbf{R}^n) \\ \|d\psi\|_{L^1(\mathbf{R}^n)} \leq 1}} \left| \int_{\mathbf{R}^n} u \, d\psi \, dx \right|.$$

Proof. This follows from Theorems A.5 and A.8. \square

Theorems A.5 and A.8 also allow to extend u by density to a linear operator from $L^1_{\#}(\mathbf{R}^n, \Lambda^k \mathbf{R}^n)$ to $\Lambda^k \mathbf{R}^n$.

Proposition 2.7. *If $u \in D_k(\mathbf{R}^n)$, then $\langle u, \varphi \rangle \in \Lambda^k \mathbf{R}^n$ is well defined for every $\varphi \in L^1_{\#}(\mathbf{R}^n, \Lambda^k \mathbf{R}^n)$.*

We shall also consider the subspace generated by continuous functions.

Definition 2.8. The space $V_k(\mathbf{R}^n)$ is the closure of the set of bounded continuous functions in $D_k(\mathbf{R}^n)$.

2.3. Characterization of $D_n(\mathbf{R}^n)$

The space $D_n(\mathbf{R}^n)$ is well known; it is $L^\infty(\mathbf{R}^n)/\mathbf{R}$.

Proposition 2.9. *The spaces $D_n(\mathbf{R}^n)$ and $L^\infty(\mathbf{R}^n)/\mathbf{R}$ are isometrically isomorphic.*

Proof. If $u \in L^\infty(\mathbf{R}^n)$, then for every $\varphi \in \mathcal{D}_{\#}(\mathbf{R}^n; \Lambda^n \mathbf{R}^n)$ and $\lambda \in \mathbf{R}$,

$$\left| \int_{\mathbf{R}^n} u \varphi \, dx \right| = \left| \int_{\mathbf{R}^n} (u - \lambda) \varphi \, dx \right| \leq \|u - \lambda\|_{L^\infty(\mathbf{R}^n)} \|\varphi\|_{L^1(\mathbf{R}^n)}.$$

Conversely, if $u \in D_n(\mathbf{R}^n)$, then

$$\varphi \mapsto \ell(\varphi) = \int_{\mathbf{R}^n} u \varphi \, dx$$

is a linear continuous mapping from $L^1_{\#}(\mathbf{R}^n, \Lambda^n \mathbf{R}^n)$ to $\Lambda^n \mathbf{R}^n \cong \mathbf{R}$. By the Hahn–Banach theorem there is an extension $\tilde{\ell}$ to $L^1(\mathbf{R}^n, \Lambda^n \mathbf{R}^n)$. This extension is represented as $\tilde{\ell}(\varphi) = \int_{\mathbf{R}^n} \tilde{u} \varphi \, dx$ with $\|\tilde{u}\|_{L^\infty(\mathbf{R}^n)} \leq \|u\|_{D_n(\mathbf{R}^n)}$. Since $\int_{\mathbf{R}^n} \tilde{u} \varphi \, dx = \int_{\mathbf{R}^n} u \varphi \, dx$ for every $\varphi \in \mathcal{D}_{\#}(\mathbf{R}^n, \Lambda^n \mathbf{R}^n)$, there is $\lambda \in \mathbf{R}$ such that $u - \lambda = \tilde{u}$. \square

2.4. Geometric characterization of $V_1(\mathbf{R}^n)$

The definition of $D_k(\mathbf{R}^n)$, and hence that of $V_k(\mathbf{R}^n)$, rely on compactly supported closed smooth forms (Definition 2.4), or equivalently on compactly supported exact forms (Proposition 2.6). Compactly supported smooth forms ensure that the definition makes sense for distributions. There is a more geometrical characterization for continuous functions, which extends by density to $V_1(\mathbf{R}^n)$ and $V_{n-1}(\mathbf{R}^n)$.

Proposition 2.10. *For every $u \in C(\mathbf{R}^n)$,*

$$\|u\|_{D_1(\mathbf{R}^n)} = \sup_{\Omega} \frac{1}{\mathcal{H}^{n-1}(\partial\Omega)} \left| \int_{\partial\Omega} u(y) \nu(y) \, d\mathcal{H}^{n-1}(y) \right|, \tag{2.1}$$

where the supremum is taken over bounded domains Ω with a smooth connected boundary, and $\nu(y)$ is the unit exterior normal vector to the boundary at $y \in \partial\Omega$.

Proof. Let Ω be a bounded domains with a smooth connected boundary. Let $\rho \in \mathcal{D}(B(0, 1))$ be such that $\rho \geq 0$ and $\int_{\mathbf{R}^n} \rho \, dx = 1$, let $\rho_\varepsilon(x) = \rho(x/\varepsilon)/\varepsilon^n$, and define

$$\varphi_\varepsilon(x) = \frac{1}{\mathcal{H}^{n-1}(\partial\Omega)} \int_{\partial\Omega} \rho_\varepsilon(y-x)\nu(y) \, d\mathcal{H}^{n-1}(y).$$

Since $\|\varphi_\varepsilon\|_1 \leq 1$ and $d\varphi_\varepsilon = 0$,

$$\left| \int_{\mathbf{R}^n} u\varphi_\varepsilon \, dx \right| \leq \|u\|_{D_1(\mathbf{R}^n)}.$$

Since u is continuous, $\rho_\varepsilon * u \rightarrow u$ as $\varepsilon \rightarrow 0$ uniformly on every compact subset of \mathbf{R}^n , and

$$\begin{aligned} \left| \int_{\mathbf{R}^n} u\varphi_\varepsilon \, dx \right| &= \frac{1}{\mathcal{H}^{n-1}(\partial\Omega)} \left| \int_{\partial\Omega} (\rho_\varepsilon * u)(y)\nu(y) \, d\mathcal{H}^{n-1}(y) \right| \\ &\rightarrow \frac{1}{\mathcal{H}^{n-1}(\partial\Omega)} \left| \int_{\partial\Omega} u(y)\nu(y) \, d\mathcal{H}^{n-1}(y) \right| \end{aligned}$$

as $\varepsilon \rightarrow 0$. Therefore

$$\frac{1}{\mathcal{H}^{n-1}(\partial\Omega)} \left| \int_{\partial\Omega} u(y)\nu(y) \, d\mathcal{H}^{n-1}(y) \right| \leq \|u\|_{D_1(\mathbf{R}^n)}.$$

Conversely, let A denote the right-hand side of (2.1). First note that for every bounded open set Ω with a smooth boundary that is not necessarily connected,

$$\left| \int_{\partial\Omega} u(y)\nu(y) \, d\mathcal{H}^{n-1}(y) \right| \leq A\mathcal{H}^{n-1}(\partial\Omega).$$

By Proposition 2.6, we need to evaluate, for $\varphi \in \mathcal{D}(\mathbf{R}^n, \Lambda^0\mathbf{R})$,

$$\left| \int_{\mathbf{R}^n} u\nabla\varphi \, dx \right| = \left| \int_{\mathbf{R}^n} \varphi\nabla u \, dx \right|.$$

One has

$$\int_{\mathbf{R}^n} \varphi\nabla u \, dx = \int_0^\infty \int_{\{x \in \mathbf{R}^n: \varphi(x) > s\}} \nabla u(x) \, dx \, ds - \int_{-\infty}^0 \int_{\{x \in \mathbf{R}^n: \varphi(x) < s\}} \nabla u(x) \, dx \, ds.$$

For every $s > 0$, the set $\{x \in \mathbf{R}^n: \varphi(x) > s\}$ is open and bounded. Moreover, by Sard’s lemma, for almost every $s > 0$, for every $y \in \varphi^{-1}(\{s\})$, $\nabla\varphi(y) \neq 0$. Hence $\partial\{x \in \mathbf{R}^n: \varphi > s\}$ is smooth and

$$\begin{aligned} \left| \int_{\{x \in \mathbf{R}^n: \varphi > s\}} \nabla u(x) \, dx \right| &= \left| \int_{\partial\{x \in \mathbf{R}^n: \varphi(x) > s\}} u(y)v(y) \, d\mathcal{H}^{n-1}(y) \right| \\ &\leq A\mathcal{H}^{n-1}(\partial\{x \in \mathbf{R}^n: \varphi(x) > s\}). \end{aligned}$$

A similar reasoning for $s < 0$, and the integration with respect to s allow to conclude with the co-area formula [9,10]:

$$\begin{aligned} \left| \int_{\mathbf{R}^n} u \nabla \varphi \, dx \right| &\leq A \int_{\mathbf{R}^+} \mathcal{H}^{n-1}(\partial\{x \in \mathbf{R}^n: \varphi(x) > s\}) + \mathcal{H}^{n-1}(\partial\{x \in \mathbf{R}^n: \varphi(x) < -s\}) \, ds \\ &= A \int_{\mathbf{R}^n} |\nabla \varphi| \, dx. \quad \square \end{aligned}$$

2.5. Geometric characterization of $V_{n-1}(\mathbf{R}^n)$

Proposition 2.11. *If $n \geq 2$, for every $u \in C(\mathbf{R}^n)$,*

$$\|u\|_{D_{n-1}(\mathbf{R}^n)} = \sup_{\gamma \in C^1(S^1; \mathbf{R}^n)} \frac{1}{\|\dot{\gamma}\|_{L^1(S^1)}} \left| \int_{S^1} u(\gamma(t)) \dot{\gamma}(t) \, dt \right|.$$

The proof repeats the argument of Bourgain and Brezis for the equivalence between the inequality (1.7) and

$$\int_{S^1} u(\gamma(t)) \dot{\gamma}(t) \, dt \leq C_{s,p} \|\dot{\gamma}\|_{L^1(S^1)} \|u\|_{W^{s,p}(\mathbf{R}^n)},$$

for every $u \in W^{s,p}(\mathbf{R}^n)$ with $sp = n$ [3].

Proof. First note that

$$\|u\|_{D_{n-1}(\mathbf{R}^n)} = \sup_{\substack{f \in \mathcal{D}(\mathbf{R}^n; \mathbf{R}^n) \\ \operatorname{div} f = 0 \\ \|f\|_{L^1(\mathbf{R}^n; \mathbf{R}^n)} \leq 1}} \left| \int_{\mathbf{R}^n} u f \, dx \right|.$$

Let $\rho \in \mathcal{D}(B(0, 1))$ be such that $\rho \geq 0$ and $\int_{\mathbf{R}^n} \rho \, dx = 1$, and let $\rho_\varepsilon(x) = \rho(x/\varepsilon)/\varepsilon^n$. Define

$$f_\varepsilon(x) = \frac{1}{\|\dot{\gamma}\|_{L^1(S^1)}} \int_{S^1} \rho(\gamma(t) - x) \dot{\gamma}(t) \, dt.$$

One has $\|f_\varepsilon\|_{L^1(\mathbf{R}^n)} \leq 1$ and $\operatorname{div} f_\varepsilon = 0$, therefore

$$\left| \int_{\mathbf{R}^n} u f_\varepsilon \, dx \right| \leq \|u\|_{D_{n-1}(\mathbf{R}^n)}.$$

The proof continues as for the first part of Proposition 2.10.

The converse inequality comes from on a result of Smirnov [18], which states that for every $R > 0$ and for every $f \in \mathcal{D}(B(0, R); \mathbf{R}^n)$ there exists $(\gamma_m^\ell)_{1 \leq m, \ell}$ in $C^1(S^1; B(0, R))$ and $(\lambda_m^\ell)_{1 \leq m, \ell}$ in \mathbf{R} such that for every $m \geq 1$,

$$\sum_{\ell \geq 1} |\lambda_m^\ell| \|\dot{\gamma}_m^\ell\|_{L^1(S^1)} \leq \|f\|_{L^1(\mathbf{R}^n)},$$

and for every $u \in C(B(0, R))$

$$\sum_{\ell=1}^{\infty} \lambda_m^\ell \int_{S^1} u(\gamma_m^\ell(t)) \dot{\gamma}_m^\ell \, dt \rightarrow \int_{\mathbf{R}^n} u f \, dx$$

as $m \rightarrow \infty$. \square

2.6. Geometric characterization of $\mathbb{V}_k(\mathbf{R}^n)$

The characterization of the seminorm $\|\cdot\|_{D_k(\mathbf{R}^n)}$ of Proposition 2.10, relies essentially on the fact that the seminorm could be evaluated by considering differential of scalar functions, while in Proposition 2.11 it relied on the decomposition result of Smirnov. Those facts do not hold anymore for $1 < k < n - 1$, but there is an equivalent geometrical seminorm expressed in terms of real polyhedral chains. Let us first recall some basic facts and notations about currents and polyhedral chains [10,17].

Definition 2.12. The space of k -dimensional currents $\mathcal{D}_k(\mathbf{R}^n)$ is the topological dual of $\mathcal{D}(\mathbf{R}^n; \Lambda^k \mathbf{R}^n)$.

The boundary of a current $T \in \mathcal{D}_k(\mathbf{R}^n)$, is $\partial T \in \mathcal{D}_{k-1}(\mathbf{R}^n)$ defined by

$$\langle \partial T, \varphi \rangle = \langle T, d\varphi \rangle.$$

The *mass* of a current is

$$\mathbf{M}(T) = \sup\{\langle T, \varphi \rangle : \varphi \in \mathcal{D}(\mathbf{R}^n; \Lambda^k \mathbf{R}^n) \text{ and } \forall x \in \mathbf{R}^n, |\varphi(x)| \leq 1\}.$$

The *support* $\operatorname{supp} T$ of a current $T \in \mathcal{D}_k(\mathbf{R}^n)$ is the complement of the largest open set U such that $\langle T, \varphi \rangle = 0$ when $\operatorname{supp} \varphi \subset U$. For every integer k , let

$$S_k = \left\{ \lambda \in \mathbf{R}^k : \sum_{i=1}^k \lambda_k \leq 1 \text{ and } \forall 1 \leq i \leq k, \lambda_i \geq 0 \right\}.$$

For every finite collection $(x_i)_{1 \leq i \leq k}$ in \mathbf{R}^n , the current $\llbracket x_0, \dots, x_k \rrbracket \in \mathcal{D}_k(\mathbf{R}^n)$ is defined by

$$\langle \llbracket x_0, \dots, x_k \rrbracket, \varphi \rangle = \int_{S_k} \left\langle \varphi \left(x_0 + \sum_{i=1}^k \lambda_i x_i \right), x_1 - x_0 \wedge \dots \wedge x_k - x_0 \right\rangle d\lambda.$$

Definition 2.13. A current $T \in \mathcal{D}_k(\mathbf{R}^n)$ is a real polyhedral chain if there is $(x_j^i)_{1 \leq i \leq m, 1 \leq j \leq k}$ in \mathbf{R}^n and $(\mu_i)_{1 \leq i \leq m}$ in \mathbf{R} such that

$$T = \sum_{i=1}^m \mu_i \llbracket x_0^i, \dots, x_k^i \rrbracket.$$

The set of k -dimensional real polyhedral chains is denoted by $\mathbf{P}_k(\mathbf{R}^n)$. Every real polyhedral chain has a compact support and a finite mass. Hence, $\langle T, u \rangle$ is well defined when $u : \mathbf{R}^n \rightarrow \Lambda^k \mathbf{R}^n$ is continuous. Moreover, if $u : \mathbf{R}^n \rightarrow \mathbf{R}$ is continuous then $\langle T, u \rangle \in (\Lambda^k \mathbf{R}^n)' \cong \Lambda^k \mathbf{R}^n$ is naturally defined.

Definition 2.14. If $u : \mathbf{R}^n \rightarrow \mathbf{R}$ is continuous, let

$$\|u\|_{\tilde{V}_k(\mathbf{R}^n)} = \sup_{\substack{P \in \mathbf{P}_{n-k}(\mathbf{R}^n) \\ \partial P = 0 \\ \mathbf{M}(P) \leq 1}} \langle P, u \rangle.$$

The seminorm $\|u\|_{\tilde{V}_k(\mathbf{R}^n)}$ measures the oscillation of the function u through its integral on k -dimensional real polyhedral chains without boundary.

Theorem 2.15. For every $n \geq 1$ and $1 \leq k \leq n - 1$, there exists $c > 0$ such that for every $u \in C(\mathbf{R}^n)$,

$$\|u\|_{\tilde{V}_k(\mathbf{R}^n)} \leq \|u\|_{\mathcal{D}_k(\mathbf{R}^n)} \leq c \|u\|_{\tilde{V}_k(\mathbf{R}^n)}.$$

Proof. Given P in $\mathbf{P}_{n-k}(\mathbf{R}^n)$, let

$$\varphi_\varepsilon(x) = \langle P, \rho_\varepsilon(\cdot - x) \rangle,$$

where $\rho_\varepsilon = \rho(\cdot/\varepsilon)/\varepsilon^n$ with $\rho \in \mathcal{D}(\mathbf{R}^n)$, $\rho \geq 0$ and $\int_{\mathbf{R}^n} \rho dx = 1$ and where $*$ denotes the Hodge duality between $\Lambda^k \mathbf{R}^n$ and $\Lambda^{n-k} \mathbf{R}^n$. One checks that $d\varphi_\varepsilon = 0$, $\|\varphi_\varepsilon\|_{L^1(\mathbf{R}^n)} \leq \mathbf{M}(P)$ and

$$\int_{\mathbf{R}^n} u \varphi_\varepsilon dx = * \langle P, \rho_\varepsilon * u \rangle.$$

Since $\rho_\varepsilon * u \rightarrow u$ uniformly as $\varepsilon \rightarrow 0$,

$$\|u\|_{V_k(\mathbf{R}^n)} \leq \|u\|_{\mathcal{D}_k(\mathbf{R}^n)}.$$

The converse inequality is based on the deformation Theorem for currents [10,17]. It states that given $T \in \mathcal{D}_k(\mathbf{R}^n)$ with $\mathbf{M}(T) + \mathbf{M}(\partial T) < \infty$, for every $\varepsilon > 0$, there exists $P \in \mathbf{P}_k(\mathbf{R}^n)$, $S \in \mathcal{D}_k(\mathbf{R}^n)$ and $R \in \mathcal{D}_{k+1}(\mathbf{R}^n)$ such that

$$T - P = \partial R + S,$$

with

$$\begin{aligned} \mathbf{M}(P) &\leq c\mathbf{M}(T), & \mathbf{M}(\partial P) &\leq c\mathbf{M}(\partial T), \\ \mathbf{M}(R) &\leq c\varepsilon\mathbf{M}(T), & \mathbf{M}(S) &\leq c\varepsilon\mathbf{M}(\partial T), \end{aligned}$$

and

$$\begin{aligned} \text{supp } P \cup \text{supp } R &\subset \{x \in \mathbf{R}^n : \text{dist}(x, \text{supp } T) < 2\sqrt{n+k}\}, \\ \text{supp } \partial P \cup \text{supp } S &\subset \{x \in \mathbf{R}^n : \text{dist}(x, \text{supp } \partial T) < 2\sqrt{n+k}\}. \end{aligned}$$

This implies that if $T \in \mathbf{P}_k(\mathbf{R}^n)$ and $U \subset \mathbf{R}^n$ is open and bounded, and $\text{supp } T \subset U$ and $\partial T = 0$, there exists a sequence $(P_\varepsilon)_{\varepsilon>0}$ in $\mathbf{P}_k(\mathbf{R}^n)$ with $\partial P_\varepsilon = 0$ and $\text{supp } P_\varepsilon \subset U$ such that for every $u \in C(\mathbf{R}^n; \mathbf{R}^n)$,

$$\mathbf{M}(P_\varepsilon) \leq c\mathbf{M}(T), \tag{2.2}$$

and

$$\langle P_\varepsilon, u \rangle \rightarrow \langle T, u \rangle \tag{2.3}$$

as $\varepsilon \rightarrow 0$.

Now given $\varphi \in \mathcal{D}_\#(\mathbf{R}^n; \Lambda^k \mathbf{R}^n)$, consider $T \in \mathcal{D}_{n-k}(\mathbf{R}^n)$ defined by

$$\langle T, v \rangle = \int_{\mathbf{R}^n} \varphi \wedge v \, dx.$$

Since T has compact support, $\partial T = 0$ and $\mathbf{M}(T) \leq \|\varphi\|_{L^1(\mathbf{R}^n)}$, there is a sequence $(P_\varepsilon)_{\varepsilon>0}$ in $\mathbf{P}_{n-k}(\mathbf{R}^n)$ such that $\partial P_\varepsilon = 0$, $\text{supp } P_\varepsilon \subset U$, (2.2) and (2.3), where U is a fixed open bounded set such that $\text{supp } T \subset U$. \square

3. Basic properties of $\mathbf{D}_k(\mathbf{R}^n)$

3.1. Mutual injections

The collection of spaces $\mathbf{D}_k(\mathbf{R}^n)$ is a decreasing sequence of spaces.

Theorem 3.1. *Let $k \leq \ell$. If $u \in \mathbf{D}_\ell(\mathbf{R}^n)$, then $u \in \mathbf{D}_k(\mathbf{R}^n)$, and*

$$\|u\|_{\mathbf{D}_k(\mathbf{R}^n)} \leq C \|u\|_{\mathbf{D}_\ell(\mathbf{R}^n)},$$

where C does not depend on u .

Proof. Let $\varphi \in \mathcal{D}_{\#}(\mathbf{R}^n; \Lambda^k \mathbf{R}^n)$. If $\alpha \in \Lambda^{\ell-k} \mathbf{R}^n$, then $\alpha \wedge \varphi \in \mathcal{D}_{\#}(\mathbf{R}^n; \Lambda^{\ell} \mathbf{R}^n)$. Therefore

$$\left| \int_{\mathbf{R}^n} u \alpha \wedge \varphi \, dx \right| \leq \|u\|_{D_{\ell}(\mathbf{R}^n)} \|\alpha \wedge \varphi\|_{L^1(\mathbf{R}^n)} \leq \|u\|_{D_{\ell}(\mathbf{R}^n)} \|\alpha\| \|\varphi\|_{L^1(\mathbf{R}^n)}.$$

Taking the supremum over $\alpha \in \Lambda^{\ell-k} \mathbf{R}^n$ with $|\alpha| \leq 1$ leads to the conclusion. \square

3.2. Extension theory

If $n < N$, functions in $D_k(\mathbf{R}^n)$ can be extended to functions in $D_k(\mathbf{R}^N)$. This extension operator is an isomorphism on its image.

Theorem 3.2. Let $u \in \mathcal{D}'(\mathbf{R}^n)$. Define $U \in \mathcal{D}'(\mathbf{R}^N)$ by

$$\int_{\mathbf{R}^N} U(z) \varphi(z) \, dz = \int_{\mathbf{R}^n} u(x) \int_{\mathbf{R}^{N-n}} \varphi(x, y) \, dy \, dx.$$

For every $1 \leq k \leq n$,

$$c \|U\|_{D_k(\mathbf{R}^N)} \leq \|u\|_{D_k(\mathbf{R}^n)} \leq C \|U\|_{D_k(\mathbf{R}^N)},$$

where $c, C > 0$ are independent of u and U .

Proof. By induction, it is sufficient to consider the case $N = n + 1$.

First let us estimate $\|U\|_{D_k(\mathbf{R}^N)}$. Consider $\Phi \in \mathcal{D}_{\#}(\mathbf{R}^N; \Lambda^k \mathbf{R}^N)$. It can be written as

$$\Phi = \Phi_0 + \Phi_1 \wedge \omega_N,$$

where $\Phi_0 \in \mathcal{D}_{\#}(\mathbf{R}^N; \Lambda^k \mathbf{R}^n)$ and $\Phi_1 \in \mathcal{D}_{\#}(\mathbf{R}^N; \Lambda^{k-1} \mathbf{R}^n)$. Define

$$\varphi(x) = \int_{\mathbf{R}} \Phi(x, t) \, dt, \quad \varphi_0(x) = \int_{\mathbf{R}} \Phi_0(x, t) \, dt, \quad \varphi_1(x) = \int_{\mathbf{R}} \Phi_1(x, t) \, dt.$$

For $m = 1, 2$,

$$d\varphi_m(x) = \sum_{i=1}^n \omega_i \wedge \frac{\partial \varphi_m}{\partial x_i} = \int_{\mathbf{R}} \sum_{i=1}^N \omega_i \wedge \frac{\partial \Phi_m}{\partial x_i} \, dt = 0$$

and

$$\int_{\mathbf{R}^n} \varphi_m \, dx = \int_{\mathbf{R}^n} \int_{\mathbf{R}} \Phi_m \, dt \, dx = 0.$$

Since $\varphi = \varphi_0 + \varphi_1 \wedge \omega_N$, one has

$$\int_{\mathbf{R}^N} U \Phi dz = \int_{\mathbf{R}^n} u \varphi dx = \int_{\mathbf{R}^n} u \varphi_0 dx + \int_{\mathbf{R}^n} u \varphi_1 dx \wedge \omega_N$$

and therefore,

$$\left| \int_{\mathbf{R}^N} U \Phi dz \right| \leq \|u\|_{D_k(\mathbf{R}^n)} \|\varphi_0\|_{L^1(\mathbf{R}^n)} + \|u\|_{D_{k-1}(\mathbf{R}^n)} \|\varphi_1\|_{L^1(\mathbf{R}^n)} |\omega_N|.$$

When $k > 1$, the conclusion comes from Theorem 3.1 and from the inequality

$$\|\varphi_0\|_{L^1(\mathbf{R}^n)} + \|\varphi_1\|_{L^1(\mathbf{R}^n)} \leq \|\Phi_0\|_{L^1(\mathbf{R}^n)} + \|\Phi_1 \wedge \omega_N\|_{L^1(\mathbf{R}^n)} \leq \|\Phi_0 + \Phi_1 \wedge \omega_N\|_{L^1(\mathbf{R}^n)}.$$

The last inequality comes from the fact that $\Phi_0(x)$ and $\Phi_1(x) \wedge \omega_N$ are orthogonal for every $x \in \mathbf{R}^N$. When $k = 1$, one has $\varphi_1 = 0$, and the conclusion comes similarly.

Conversely, let us now estimate $\|u\|_{D_k(\mathbf{R}^n)}$ by Proposition 2.6. Let $\psi \in \mathcal{D}(\mathbf{R}^n; \Lambda^{k-1}\mathbf{R}^n)$. Consider a family $(\eta_\lambda)_{\lambda>0}$ in $\mathcal{D}(\mathbf{R})$ such that $\eta_\lambda \geq 0$, $\int_{\mathbf{R}} \eta_\lambda dt = 1$ and $\int_{\mathbf{R}} |\eta'_\lambda| dt \rightarrow 0$ as $\lambda \rightarrow \infty$, and let $\Psi_\lambda(x, t) = \eta_\lambda(t)\psi(x)$. For every $\lambda > 0$,

$$\int_{\mathbf{R}^N} U d\Psi_\lambda dz = \int_{\mathbf{R}^n} U (d\eta_\lambda \wedge \psi + \eta_\lambda d\psi) dz = 0 + \int_{\mathbf{R}^n} u d\psi dx.$$

Therefore,

$$\left| \int_{\mathbf{R}^n} u d\psi dx \right| = \left| \int_{\mathbf{R}^N} U d\Psi_\lambda dz \right| \leq \|U\|_{D_k(\mathbf{R}^N)} (\|\psi\|_{L^1(\mathbf{R}^n)} + \|d\psi\|_{L^1(\mathbf{R}^n)} \|\eta'_\lambda\|_{L^1(\mathbf{R})}).$$

Letting $\lambda \rightarrow \infty$ yields the conclusion. \square

Remark 3.3. When $k < n$, the result of Theorem 3.2 is optimal: there are functions in $D_k(\mathbf{R}^n)$ whose extension to \mathbf{R}^N does not belong to $D_K(\mathbf{R}^N)$ for $K > k$ (see Proposition 4.6). On the other hand, the extension of a function in $D_n(\mathbf{R}^n)$ lies in $D_N(\mathbf{R}^N)$ by Proposition 2.9. In view of the trace theory of the next section, one could wonder whether when $1 \leq k < n$, there is *another* extension in $D_K(\mathbf{R}^N)$ with $K > k$.

3.3. Trace theory

The restriction of continuous functions from \mathbf{R}^N to \mathbf{R}^n can be extended to a continuous operator from $V_K(\mathbf{R}^N)$ to $V_k(\mathbf{R}^n)$ when $N - K = n - k$.

Theorem 3.4. *Let $n \leq N$, $1 \leq K \leq N$ and $k = K - (N - n)$. Let $U \in V_k(\mathbf{R}^N)$ be continuous. Define for $x \in \mathbf{R}^n$,*

$$u(x) = U(x, 0).$$

Then $u \in V_k(\mathbf{R}^n)$, and

$$\|u\|_{D_k(\mathbf{R}^n)} \leq \|U\|_{D_k(\mathbf{R}^N)}.$$

Proof. By induction, we can assume $N = n + 1$. Let $\varphi \in \mathcal{D}_\#(\mathbf{R}^n; \Lambda^k \mathbf{R}^n)$, let $\rho \in \mathcal{D}(\mathbf{R})$ such that $\rho \geq 0$ and $\int_{\mathbf{R}} \rho dt = 1$ and let $\rho_\varepsilon(t) = \rho(t/\varepsilon)/\varepsilon$. Let $\Phi_\varepsilon(x, t) = \rho_\varepsilon(t)\psi(x) \wedge \omega_N$. Since $\Phi_\varepsilon \in \mathcal{D}_\#(\mathbf{R}^N; \Lambda^k \mathbf{R}^N)$ and u is continuous,

$$\begin{aligned} \left| \int_{\mathbf{R}^n} u\varphi dx \right| &= \lim_{\varepsilon \rightarrow 0} \left| \int_{\mathbf{R}^n} \int_{\mathbf{R}} u\Phi_\varepsilon dt dx \right| \\ &\leq \|U\|_{D_k(\mathbf{R}^N)} \lim_{\varepsilon \rightarrow 0} \|\Phi_\varepsilon\|_{L^1(\mathbf{R}^N)} \leq \|U\|_{D_k(\mathbf{R}^N)} \|\varphi\|_{L^1(\mathbf{R}^n)}. \quad \square \end{aligned}$$

4. Examples of functions in $D_k(\mathbf{R}^n)$

4.1. Sobolev spaces

The first class of functions in the space $D_k(\mathbf{R}^n)$ are functions in critical Sobolev spaces, which motivated the definition.

Theorem 4.1. (Bourgain and Brezis [3]) *If $u \in W^{s,p}(\mathbf{R}^n)$, $p > 1$ and $sp = n$, then for every $1 \leq k \leq n - 1$, $u \in D_k(\mathbf{R}^n)$, and*

$$\|u\|_{D_k(\mathbf{R}^n)} \leq C_{k,s,p} \|u\|_{W^{s,p}(\mathbf{R}^n)}.$$

The seminorm on the right-hand side is the Sobolev semi-norm. For $0 < s < 1$ it is defined as

$$\|u\|_{W^{s,p}(\mathbf{R}^n)}^p = \int_{\mathbf{R}^n \times \mathbf{R}^n} \frac{|u(x) - u(y)|^p}{|x - y|^{n+sp}} dx dy.$$

Proof. This follows from the inequality

$$\left| \int_{\mathbf{R}^n} u\varphi dx \right| \leq C_{s,p} \|u\|_{W^{s,p}} \|\varphi\|_{L^1(\mathbf{R}^n)}$$

for every $\varphi \in \mathcal{D}_\#(\mathbf{R}^n; \Lambda^k \mathbf{R}^n)$ and $u \in W^{s,p}(\mathbf{R}^n)$ [3,23]. \square

Remark 4.2. The Besov spaces $B_q^{s,p}(\mathbf{R}^n)$ and the Triebel–Lizorkin spaces $F_q^{s,p}(\mathbf{R}^n)$, with $sp = n$ and $1 < q < \infty$, are embedded in $D_k(\mathbf{R}^n)$ for $1 \leq k \leq n$. This follows from the standard embeddings of these spaces in

$$W^{s',p'}(\mathbf{R}^n) = B_{p'}^{s',p'}(\mathbf{R}^n) = F_{p'}^{s',p'}(\mathbf{R}^n),$$

when $0 < s' < \min(s, 1)$, $p' \geq q$ and $s'p' = n$ [14,21].

4.2. *Locally Lipschitz functions in $\mathbf{R}^n \setminus \{0\}$*

The space $D_{n-1}(\mathbf{R}^n)$ is larger than critical Sobolev spaces. There is a simple condition for locally Lipschitz functions in $\mathbf{R}^n \setminus \{0\}$ to be in $D_{n-1}(\mathbf{R}^n)$, which is satisfied e.g. by the function $\log |x|$.

Proposition 4.3. *Let $n \geq 2$ and $u \in W_{loc}^{1,1}(\mathbf{R}^n \setminus \{0\})$. If $|x|\nabla u \in L^\infty(\mathbf{R}^n)$, then $u \in D_{n-1}(\mathbf{R}^n)$ and*

$$\|u\|_{D_{n-1}(\mathbf{R}^n)} \leq \| |x|\nabla u \|_{L^\infty(\mathbf{R}^n)}.$$

Remark 4.4. In general, $u \notin D_n(\mathbf{R}^n)$ as shows the function $u(x) = \log |x|$.

Proof. Let $f \in \mathcal{D}(\mathbf{R}^n \setminus \{0\}; \mathbf{R}^n)$ be such that $\operatorname{div} f = 0$. Lemma 4.5 yields

$$\left| \int_{\mathbf{R}^n} u f \, dx \right| = \left| \int_{\mathbf{R}^n} x(\nabla u \cdot f) \, dx \right| \leq \| \nabla u |x| \|_{L^\infty(\mathbf{R}^n)} \| f \|_{L^1(\mathbf{R}^n)}.$$

Therefore, for every $\varphi \in \mathcal{D}_\#(\mathbf{R}^n \setminus \{0\}; \mathbf{R}^n)$,

$$\left| \int_{\mathbf{R}^n} u \varphi \, dx \right| \leq \| \nabla u |x| \|_{L^\infty(\mathbf{R}^n)} \| \varphi \|_{L^1(\mathbf{R}^n)}.$$

Since $\{0\}$ has vanishing n -capacity (Lemma A.2),

$$\mathcal{D}_\#(\mathbf{R}^n \setminus \{0\}; \Lambda^{n-1}\mathbf{R}) \text{ is dense in } \mathcal{D}_\#(\mathbf{R}^n; \Lambda^{n-1}\mathbf{R})$$

(Theorem A.5). This concludes the proof. \square

Lemma 4.5. *Let $u \in W_{loc}^{1,1}(\mathbf{R}^n \setminus \{0\})$ and $f \in \mathcal{D}(\mathbf{R}^n \setminus \{0\}; \mathbf{R}^n)$. If $\operatorname{div} f = 0$, then*

$$\int_{\mathbf{R}^n} u f \, dx = - \int_{\mathbf{R}^n} x(f \cdot \nabla u) \, dx.$$

Proof. By integration by parts,

$$\int_{\mathbf{R}^n} x(f \cdot \nabla u) \, dx = - \int_{\mathbf{R}^n} x(\operatorname{div} f)u \, dx - \int_{\mathbf{R}^n} u f \, dx.$$

The conclusion comes from the assumption $\operatorname{div} f = 0$. \square

4.3. *Examples of functions in $D_k(\mathbf{R}^n)$*

Proposition 4.3 and Theorem 3.2 yield examples of functions in the spaces $D_k(\mathbf{R}^n)$ showing that these spaces are distinct.

Proposition 4.6. *If $1 \leq \ell \leq n$, then*

$$\log\left(\sum_{i=1}^{\ell} |x_i|^2\right) \in D_k(\mathbf{R}^n)$$

if and only if $1 \leq k < \ell$.

Remark 4.7. An immediate consequence of Proposition 4.6 is that the spaces $D_k(\mathbf{R}^n)$ are all different.

Remark 4.8. Proposition 4.6 is consistent with the isomorphism $D_n(\mathbf{R}^n) \cong L^\infty(\mathbf{R}^n)/\mathbf{R}$ (Proposition 2.9).

Proof of Proposition 4.6. For every $x \in \mathbf{R}^n \setminus \{0\}$, $|x| |\nabla(\log|x|)| = 1$. Therefore, for $\ell \geq 2$, by Proposition 4.3,

$$\log\left(\sum_{i=1}^{\ell} |x_i|^2\right) \in D_{\ell-1}(\mathbf{R}^\ell).$$

By Theorem 3.2, this remains valid for the extension to \mathbf{R}^n :

$$\log\left(\sum_{i=1}^{\ell} |x_i|^2\right) \in D_{\ell-1}(\mathbf{R}^n).$$

Hence, by Theorem 3.1, if $1 \leq k < \ell$,

$$\log\left(\sum_{i=1}^{\ell} |x_i|^2\right) \in D_k(\mathbf{R}^n).$$

On the other hand, suppose for contradiction that, for some $k \geq \ell$,

$$\log\left(\sum_{i=1}^{\ell} |x_i|^2\right) \in D_k(\mathbf{R}^n).$$

By Theorem 3.1, this would be true for $k = \ell$. By Theorem 3.2,

$$\log\left(\sum_{i=1}^{\ell} |x_i|^2\right) \in D_\ell(\mathbf{R}^\ell) = L^\infty(\mathbf{R}^\ell)/\mathbf{R},$$

which is absurd. \square

Proposition 4.6 includes as a special case estimates obtained by Bourgain and Brezis:

Corollary 4.9. (Bourgain and Brezis [3]) *Let $f \in L^1(\mathbf{R}^2; \mathbf{R}^2)$. If $\operatorname{div} f = 0$ in the sense of distributions, then*

$$\log \frac{1}{|x|} * f \in L^\infty(\mathbf{R}^2; \mathbf{R}^2).$$

Other interesting examples can be obtained in a similar way:

Proposition 4.10. *If $1 \leq \ell \leq n$ and $0 < \alpha \leq 1$, then*

$$\left(\left| \log \left(\sum_{i=1}^{\ell} |x_i|^2 \right) \right| + 1 \right)^\alpha \in D_k(\mathbf{R}^n)$$

if and only if $1 \leq k < \ell$.

Note that

$$\left(\left| \log \left(\sum_{i=1}^{\ell} |x_i|^2 \right) \right| + 1 \right)^\alpha \notin W_{\text{loc}}^{s,p}(\mathbf{R}^n)$$

when $\ell < n$ (otherwise it would be continuous on almost every hyperplane). Hence the examples provided here do not belong to critical Sobolev spaces.

5. Relation with BMO(\mathbf{R}^n)

5.1. Preliminaries

Let us first recall some facts about the space of functions with bounded mean oscillation BMO(\mathbf{R}^n) [20].

The space BMO(\mathbf{R}^n) is defined as the set of functions $u \in L^1_{\text{loc}}(\mathbf{R}^n)$ such that

$$\|u\|_{\text{BMO}(\mathbf{R}^n)} = \sup_B \frac{1}{\mathcal{L}^n(B)^2} \int_B \int_B |u(x) - u(y)| \, dx \, dy < \infty,$$

where the supremum is taken on balls $B \subset \mathbf{R}^n$.

The space BMO(\mathbf{R}^n) is the dual space of the real Hardy space $H^1(\mathbf{R}^n)$, which can be characterized as the space of functions $f \in L^1(\mathbf{R}^n)$ such that $\mathcal{R}_i f \in L^1(\mathbf{R}^n)$, where \mathcal{R}_i denotes the Riesz transform (defined by $\mathcal{R}_i f = x_i/|x|^{n+1} * f$ when $f \in \mathcal{D}(\mathbf{R}^n)$). One can take

$$\|f\|_{H^1(\mathbf{R}^n)} = \|f\|_{L^1(\mathbf{R}^n)} + \sum_{i=1}^n \|\mathcal{R}_i f\|_{L^1(\mathbf{R}^n)}.$$

If K_n denotes the fundamental solution of the Laplacian $\Delta = \sum_{i=1}^n \partial^2/\partial x_i^2$ (i.e. K_n is Newton’s kernel), and $f \in H^1(\mathbf{R}^n)$, one has:

$$\frac{\partial^2}{\partial x_i \partial x_j} (K_n * f) \in L^1(\mathbf{R}^n)$$

and

$$\left\| \frac{\partial^2}{\partial x_i \partial x_j} (K_n * f) \right\|_{L^1(\mathbf{R}^n)} \leq C \|f\|_{H^1(\mathbf{R}^n)}.$$

Finally, $\mathcal{D}(\mathbf{R}^n) \cap H^1(\mathbf{R}^n)$ is dense in $H^1(\mathbf{R}^n)$.

The space of functions with vanishing mean oscillations $VMO(\mathbf{R}^n; \mathbf{R}^n)$ is the closed subspace of $VMO(\mathbf{R}^n)$ that is characterized by

$$\lim_{\varepsilon \rightarrow 0} \sup_{\mathcal{L}^n(B) \leq \varepsilon} \frac{1}{\mathcal{L}^n(B)^2} \int_B \int_B |u(x) - u(y)| \, dx \, dy = 0,$$

where the supremum is taken over balls $B \subset \mathbf{R}^n$. The critical Sobolev spaces $W^{s,p}(\mathbf{R}^n)$ with $sp = n$ are embedded in $VMO(\mathbf{R}^n)$.

Going back to the examples of the preceding section, for every $\ell \geq 1$,

$$\log \left(\sum_{i=1}^{\ell} |x_i|^2 \right) \in BMO(\mathbf{R}^n),$$

but it does not belong to $VMO(\mathbf{R}^n)$, while for $0 < \alpha < 1$ and $\ell \geq 1$,

$$\left(\left| \log \left(\sum_{i=1}^{\ell} |x_i|^2 \right) \right| + 1 \right)^{\alpha} \in VMO(\mathbf{R}^n).$$

(It is still in $BMO(\mathbf{R}^n)$ when $\alpha = 1$.) Comparing with Proposition 4.10, this gives a first insight on the relationship between $D_k(\mathbf{R}^n)$, $VMO(\mathbf{R}^n)$ and $BMO(\mathbf{R}^n)$.

Proposition 5.1. *For every $1 \leq k \leq n$, the space $VMO(\mathbf{R}^n)$ is not embedded in $D_k(\mathbf{R}^n)$.*

Remark 5.2. There is thus no inequality

$$\left| \int_{\mathbf{R}^n} u \varphi \, dx \right| \leq C \|u\|_{BMO(\mathbf{R}^n)} \|\varphi\|_{L^1(\mathbf{R}^n)},$$

for $u \in VMO(\mathbf{R}^n)$. This was remarked indirectly by Bethuel et al. [2].

5.2. Embedding in BMO

The spaces $D_k(\mathbf{R}^n)$ do not contain $BMO(\mathbf{R}^n)$ nor $VMO(\mathbf{R}^n)$. Since $L^\infty(\mathbf{R}^n)/\mathbf{R} \subset D_k(\mathbf{R}^n)$ for $1 \leq k \leq n$, they are not contained in $VMO(\mathbf{R}^n)$, but there is an embedding of $D_k(\mathbf{R}^n)$ in $BMO(\mathbf{R}^n)$:

Theorem 5.3. *Let $1 \leq k \leq n$. If $u \in D_k(\mathbf{R}^n)$, then $u \in BMO(\mathbf{R}^n)$, and*

$$\|u\|_{BMO(\mathbf{R}^n)} \leq C \|u\|_{D_k(\mathbf{R}^n)},$$

where C is independent of u .

Proof. By Theorem 3.1, we can assume $k = 1$. The seminorm of u in $\text{BMO}(\mathbf{R}^n)$ will be estimated by duality with the Hardy space $H^1(\mathbf{R}^n)$.

Let $f \in \mathcal{D}(\mathbf{R}^n) \cap H^1(\mathbf{R}^n)$. Let

$$\varphi_i = \sum_{j=1}^n \frac{\partial^2}{\partial x_i \partial x_j} (K_N * f) \omega_j.$$

Note $\varphi_j \in L^1_{\#}(\mathbf{R}^n; \Lambda^1 \mathbf{R}^n)$. Moreover,

$$\begin{aligned} \left| \int_{\mathbf{R}^n} f u \, dx \right| &\leq \sum_{i=1}^n \left| \int_{\mathbf{R}^n} \partial^2 f / \partial x_i^2 u \, dx \right| \leq \sum_{i=1}^n \left| \int_{\mathbf{R}^n} \varphi_i u \, dx \right| \\ &\leq \|u\|_{D_1(\mathbf{R}^n)} \sum_{i=1}^n \|\varphi_i\|_{L^1(\mathbf{R}^n)} \leq C \|u\|_{D_1(\mathbf{R}^n)} \|f\|_{H^1(\mathbf{R}^n)}. \end{aligned} \tag{5.1}$$

(Note that Proposition 2.7 about the well-definiteness of the duality product between $D_1(\mathbf{R}^n)$ and $L^1_{\#}(\mathbf{R}^n; \Lambda^1 \mathbf{R}^n)$ was used.) \square

Remark 5.4. A similar argument shows that for $1 < p < \infty$

$$\|u\|_{L^p(\mathbf{R}^n)} \leq \sup_{\substack{\varphi \in \mathcal{D}_{\#}(\mathbf{R}^n; \mathbf{R}^n) \\ \|\varphi\|_{L^p(\mathbf{R}^n)} \leq 1}} \left| \int_{\mathbf{R}^n} u \varphi \, dx \right|.$$

The extension of the spaces $D_k(\mathbf{R}^n)$ to the case $1 < p < \infty$ would thus not be interesting.

5.3. Completeness of $D_k(\mathbf{R}^n)$

The injection of $D_k(\mathbf{R}^n)$ provides an easy proof of the completeness of $D_k(\mathbf{R}^n)$.

Theorem 5.5. *For $1 \leq k \leq n$, the space $D_k(\mathbf{R}^n)$ is a complete Banach space modulo constants.*

Proof. Let $(u_m)_{m \geq 1}$ be a Cauchy sequence in $D_k(\mathbf{R}^n)$: $\lim_{m, l \rightarrow \infty} \|u_m - u_l\|_{D_k(\mathbf{R}^n)} = 0$. By Theorem 5.3, $(u_m)_{m \geq 1}$ is a Cauchy sequence in the Banach space $\text{BMO}(\mathbf{R}^n)$ and has thus a limit $u \in \text{BMO}(\mathbf{R}^n)$. For every $\varphi \in \mathcal{D}_{\#}(\mathbf{R}^n; \Lambda^k \mathbf{R}^n)$ and $m \geq 1$

$$\left| \int_{\mathbf{R}^n} (u_m - u) \varphi \, dx \right| = \lim_{l \rightarrow \infty} \left| \int_{\mathbf{R}^n} (u_m - u_l) \varphi \, dx \right| \leq \lim_{l \rightarrow \infty} \|u_m - u_l\|_{D_k(\mathbf{R}^n)} \|\varphi\|_{L^1(\mathbf{R}^n)}.$$

Taking the supremum over $\varphi \in \mathcal{D}_{\#}(\mathbf{R}^n; \Lambda^k \mathbf{R}^n)$ with $\|\varphi\|_{L^1(\mathbf{R}^n)} \leq 1$, one obtains $u \in D_k(\mathbf{R}^n)$ and $u_m \rightarrow u$ in $D_k(\mathbf{R}^n)$ as $m \rightarrow \infty$.

Finally, if $u \in D_k(\mathbf{R}^n)$ then, by Theorem 5.3, $\|u\|_{\text{BMO}(\mathbf{R}^n)} = 0$, and therefore u is constant. \square

5.4. Integrability of functions in $D_k(\mathbf{R}^n)$

If $u \in \text{BMO}(\mathbf{R}^n)$, $\|u\|_{\text{BMO}(\mathbf{R}^n)} \leq 1$ and $\int_{B(0,1)} u \, dx = 0$, the John–Nirenberg theorem states that

$$\int_{B(0,1)} \exp(\mu|u|) \, dx \leq c, \tag{5.2}$$

where $\mu > 0$ and $c > 0$ can be chosen independently of u . Since the spaces $D_k(\mathbf{R}^n)$ are embedded in $\text{BMO}(\mathbf{R}^n)$, this might be improved on $D_k(\mathbf{R}^n)$.

On the other hand, if $sp = n$, $0 < s < 1$, $u \in \text{W}^{s,p}(\mathbf{R}^n)$, $\|u\|_{\text{W}^{s,p}(\mathbf{R}^n)} \leq 1$ and $\int_{B(0,1)} f \, dx = 0$, then

$$\int_{B(0,1)} \exp(\mu|u|^{p/(p-1)}) \, dx \leq c, \tag{5.3}$$

where the constants c and μ are independent of u [12]. The exponent $p/(p - 1)$ cannot be improved, since $\log(1/|x|)_+^\alpha \in \text{W}^{s,p}(\mathbf{R}^n)$ when $\alpha < p/(p - 1)$ [14, p. 47]. There is not much room between (5.2) and (5.3), and it is therefore not surprising that (5.2) is optimal also in $D_k(\mathbf{R}^n)$.

Proposition 5.6. *Suppose that for every compactly supported $u \in C(\mathbf{R}^n)$, if $\|u\|_{D_k(\mathbf{R}^n)} \leq 1$ and $\int_{B(0,1)} u = 0$,*

$$\int_{B(0,1)} F(u) \, dx < \infty.$$

Then there exists $\lambda > 0$ such that

$$\int_0^\infty \Phi(s)e^{-\lambda s} \, ds < \infty.$$

Proof. Choose $y \in \mathbf{R}^n$ with $|y| = 1/2$. Consider the functions $(u_m)_{m \geq 1}$ defined by

$$u_m(x) = \min(m, (\log(2|x - y|))_-) - \min(m, (\log(2|x + y|))_-).$$

By Proposition 4.3, $\|u_m\|_{D_k(\mathbf{R}^n)}$ is bounded uniformly in m . Choose $\alpha > 0$ such that $\|\alpha u_m\|_{D_k(\mathbf{R}^n)} \leq 1$. On the other hand,

$$\begin{aligned} \int_{B(0,1)} \Phi(\alpha u_m) \, dx &= 2^{1-n} \int_{B(0,1)} \Phi(\alpha \max(\log(1/|x|), m)) \, dx = K \int_1^{e^{-m}} \Phi(-\alpha \log r) r^{n-1} \, dr \\ &= \frac{K}{\alpha} \int_0^{\alpha m} \Phi(s)e^{-ns/\alpha} \, ds \leq c. \end{aligned}$$

Letting $m \rightarrow \infty$ yields the conclusion with $\lambda = n/\alpha$. \square

The improvement of $D_k(\mathbf{R}^n)$ on $BMO(\mathbf{R}^n)$ should thus not be seen as an improvement of the integrand, but as an improvement on the domains of integration: by the trace Theorem 3.4, the embedding Theorem 5.3, and the John–Nirenberg inequality, functions in $V_k(\mathbf{R}^n)$ are exponentially integrable on $(n - k + 1)$ -dimensional subspaces.

6. Further problems

6.1. Traces of $V_1(\mathbf{R}^n)$ on $VMO(\mathbf{R}^{n-1})$

By Theorems 3.4 and 5.3, functions in $V_k(\mathbf{R}^n)$ have VMO traces on $(n - k + 1)$ -dimensional spaces. The dimension $n - k$ seems more natural: functions in $V_n(\mathbf{R}^n)$ are continuous, and hence have traces on 0-dimensional spaces, i.e. points. If there was such a trace inequality, one could define $D_0(\mathbf{R}^n) = BMO(\mathbf{R}^n)$. This notation would be consistent with the mutual injection Theorem 3.1, the extension Theorem 3.2 and the examples of Proposition 4.6. It would then be nice to have a definition of $D_k(\mathbf{R}^n)$ which encompasses the case $k = 0$. The two-dimensional case would already solve the problem of traces of $V_{n-1}(\mathbf{R}^n)$ on lines.

6.2. Geometric characterizations

By Propositions 2.10 and 2.11, the spaces $V_1(\mathbf{R}^n)$ and $V_{n-1}(\mathbf{R}^n)$ can be defined by oscillations respectively along boundaries of bounded domains and along closed curves. Further refinements would restrict the set of domains and of curves. The most striking result would be if the oscillation could be simply evaluated respectively on spheres and on circles.

The spaces $V_k(\mathbf{R}^n)$ for $1 < k < n - 1$ do not have such a simple characterization. Proposition 2.14 gives an equivalent seminorm, obtained by integration on closed real polyhedral chains without boundary. This result needs to be improved by restricting the class of sets on which the oscillations are computed for example to integer polyhedral chains without boundary, or to embedded oriented $(n - k)$ -dimensional manifolds without boundary.

6.3. Closure of the space of continuous functions

The two equivalent definitions of $VMO(\mathbf{R}^n)$ [15] suggest the definition of the closure of the bounded uniformly-continuous functions $U_k(\mathbf{R}^n)$ and of

$$W_k(\mathbf{R}^n) = \{u \in D_k(\mathbf{R}^n) : N_k(u) = 0\},$$

where

$$N_k(u) = \lim_{\varepsilon \rightarrow 0} \sup_{\substack{\varphi \in \mathcal{D}_{\#}(B(x,r); \Lambda^k \mathbf{R}^n) \\ \|\varphi\|_{L^1(\mathbf{R}^n)} \leq 1 \\ x \in \mathbf{R}^n \\ r \leq \varepsilon}} \left| \int_{B(x,r)} u \varphi \, dx \right|.$$

The space $U_k(\mathbf{R}^n)$ is contained in none of the spaces $V_k(\mathbf{R}^n)$, $W_k(\mathbf{R}^n)$ and $VMO(\mathbf{R}^n)$, while $V_k(\mathbf{R}^n)$ is not contained in $VMO(\mathbf{R}^n)$. The remaining problems are thus the understanding of the mutual relationship between $W_k(\mathbf{R}^n)$ and $VMO(\mathbf{R}^n) \cap D_k(\mathbf{R}^n)$ and of their possible embeddings in $V_k(\mathbf{R}^n)$ and in $U_k(\mathbf{R}^n)$.

6.4. *Similar spaces $E_k(\mathbf{R}^n)$*

It would have been possible to define another family of spaces with properties similar to $D_k(\mathbf{R}^n)$. For $1 \leq k \leq n$, let

$$\mathcal{D}_{\#,k}(\mathbf{R}^n; \mathbf{R}^n) = \{ \varphi \in \mathcal{D}(\mathbf{R}^n; \mathbf{R}^n) : \operatorname{div} \varphi = 0 \}$$

and the dimension of the range of φ is at most k }.

(The set $\mathcal{D}_{\#,k}(\mathbf{R}^n; \mathbf{R}^n)$ is not a vector space when $k < n$.) Define, for $1 \leq k \leq n - 1$,

$$\|u\|_{E_k(\mathbf{R}^n)} = \sup_{\substack{\varphi \in \mathcal{D}_{\#,k+1}(\mathbf{R}^n; \mathbf{R}^n) \\ \|\varphi\|_{L^1(\mathbf{R}^n)} \leq 1}} \left| \int_{\mathbf{R}^n} u \varphi \, dx \right|.$$

For $1 \leq k \leq n - 1$, the space $E_k(\mathbf{R}^n)$ is continuously embedded in $D_k(\mathbf{R}^n)$, and $E_{n-1}(\mathbf{R}^n)$ is isomorphic to $D_{n-1}(\mathbf{R}^n)$. Therefore, the critical Sobolev spaces are embedded in $E_k(\mathbf{R}^n)$ which in turn are embedded in $BMO(\mathbf{R}^n)$. Moreover, the trace property of Theorem 3.4 holds for the closure of continuous function in $E_k(\mathbf{R}^n)$. Continuous functions in $E_k(\mathbf{R}^n)$ can be characterized by Proposition 2.11 provided one restricts in the supremum $\gamma(S^1)$ to be contained in a $(k + 1)$ -dimensional affine plane. The question is about where $E_k(\mathbf{R}^n)$ lies between the spaces $D_k(\mathbf{R}^n)$ or to $D_{n-1}(\mathbf{R}^n)$. In particular, is it isomorphic to one of those?

6.5. *Composition of functions in $D_k(\mathbf{R}^n)$*

Let $F : \mathbf{R} \rightarrow \mathbf{R}$ be a Lipschitz function. If $u \in W^{s,p}(\mathbf{R}^n)$ with $0 < s \leq 1$ and $sp = n$, then $F \circ u \in W^{s,p}(\mathbf{R}^n)$ [24]. If $u \in BMO(\mathbf{R}^n)$, then $F \circ u \in BMO(\mathbf{R}^n)$ [20]. Is $F \circ u \in D_k(\mathbf{R}^n)$ when F is a Lipschitz smooth function? A simpler problem is whether $|u| \in D_k(\mathbf{R}^n)$ whenever $u \in D_k(\mathbf{R}^n)$?

If $F : \mathbf{R}^n \rightarrow \mathbf{R}^n$ is smooth, the mapping $u \mapsto u \circ F$ is a continuous operator on $W^{1,n}(\mathbf{R}^n)$ and $BMO(\mathbf{R}^n)$ if and only if F is quasiconformal [13], i.e.

$$\sup_{x \in \mathbf{R}^n} \frac{|DF(x)|^n}{|\det DF(x)|} < \infty.$$

Is it true that $u \mapsto u \circ F$ is continuous from $D_k(\mathbf{R}^n)$ to $D_k(\mathbf{R}^n)$ if F is quasiconformal, or even for F smooth and bilipschitzian?

6.6. *Localization of $D_k(\mathbf{R}^n)$*

There should be localized versions of the spaces $D_k(\mathbf{R}^n)$. There are two different definitions, depending on whether the supremum

$$\sup_{\|\varphi\|_{L^1(\mathbf{R}^n)} \leq 1} \left| \int_{\Omega} u \varphi \, dx \right|$$

is taken over smooth closed forms on Ω or over smooth closed forms on \mathbf{R}^n with support in Ω . They contain respectively $W_0^{1,n}(\Omega)$ and $W^{1,n}(\Omega)$ [7]. It would be natural for these spaces to be

embedded respectively in $\text{bmo}_z(\bar{\Omega})$ (functions whose extension by 0 to \mathbf{R}^n is in $\text{BMO}(\mathbf{R}^n)$) and the second in the larger space $\text{bmo}_r(\bar{\Omega})$ (restrictions of functions in $\text{BMO}(\mathbf{R}^n)$ to Ω) (see [8] for the definitions).

Acknowledgments

The author thanks Haïm Brezis who suggested the problem, and who encouraged and discussed the progress of this work. He also thanks Thierry De Pauw for discussions, and acknowledges the hospitality of the Laboratoire Jacques-Louis Lions, Université Pierre et Marie Curie in Paris, at which this work was initiated.

Appendix A. Density of compactly supported forms

A.1. The closure of closed k -forms

This appendix is devoted to the study of dense sets in the space $L^1(\mathbf{R}^n; \Lambda^k \mathbf{R}^n)$ of summable closed forms.

Definition A.1. Let $1 \leq p < \infty$ and $\Omega \subset \mathbf{R}^n$ be open and bounded. The p -capacity of a compact set $\Sigma \subset \Omega$ is defined as

$$\text{capa}_p(\Sigma, \Omega) = \inf \left\{ \int_{\Omega} |\nabla \eta|^p: \eta \in \mathcal{D}(\Omega), \eta = 1 \text{ in a neighborhood of } \Sigma \right\}.$$

In general $\text{capa}_p(\Sigma, \Omega)$ depends on Ω , but if $\text{capa}_p(\Sigma, \Omega) = 0$ for some bounded open set Ω , then $\text{capa}_p(\Sigma, \Omega') = 0$ for every bounded open set $\Omega' \supset \Sigma$. Therefore it makes sense to speak of sets with vanishing p -capacity without specifying Ω .

Lemma A.2. *If $n \geq 2$, for every $a \in \mathbf{R}^n$, the set $\{a\}$ has vanishing n -capacity.*

Proof. Consider the sequence

$$\eta_m(x) = \theta(\log(1/|x - a|)/m),$$

with $\theta \in C^\infty(\mathbf{R})$ and $\text{supp } \theta' \subset (0, 1)$, $\theta(0) = 0$ and $\theta(1) = 1$. \square

Remark A.3. More generally, if the 0-dimensional Hausdorff measure of Σ vanishes, then $\mathcal{H}^0(\Sigma) = 0$. Conversely, if $\text{capa}_n(\Sigma) = 0$, then for every $s > 0$, the s -dimensional Hausdorff measure of Σ vanishes [9].

In a similar way, one can prove

Lemma A.4. *There exists a sequence $(\zeta_m)_m$ in $\mathcal{D}(\mathbf{R}^n)$ such that $0 \leq \zeta_m \leq 1$, $\zeta_m \rightarrow 1$ almost everywhere and*

$$\int_{\mathbf{R}^n} |\nabla \zeta_m|^n dx \rightarrow 0$$

as $m \rightarrow \infty$.

Theorem A.5. *If $\Sigma \subset \mathbf{R}^n$ is compact and has vanishing n -capacity, then $d(\mathcal{D}(\mathbf{R}^n \setminus \Sigma; \Lambda^{k-1}\mathbf{R}^n))$ is dense in $L^1_{\#}(\mathbf{R}^n; \Lambda^k\mathbf{R}^n)$.*

The proof makes use of a result of Bourgain and Brezis.

Theorem A.6. (Bourgain and Brezis [3]) *Let $1 \leq k \leq n - 1$. For every $\varphi \in L^1_{\#}(\mathbf{R}^n; \Lambda^k\mathbf{R}^n)$, there exists $\psi \in L^{n/(n-1)}(\mathbf{R}^n; \Lambda^{k-1}\mathbf{R}^n)$ such that*

$$\begin{cases} d\psi = \varphi, \\ \delta\psi = 0. \end{cases}$$

Here δ denotes the codifferential, i.e. the adjoint of d with respect to Hodge star. This result is based on inequality (1.7). When $k = 1$, the meaningless condition $\delta\psi = 0$ is dropped and this is equivalent with the Nirenberg–Sobolev embedding.

Proof of Theorem A.5. Since the exterior differential d commutes with translations, by classical smoothing arguments, $(C^\infty \cap L^1_{\#})(\mathbf{R}^n; \Lambda^k\mathbf{R}^n)$ is dense in $L^1_{\#}(\mathbf{R}^n; \Lambda^k\mathbf{R}^n)$.

Let $\varphi \in (C^\infty \cap L^1_{\#})(\mathbf{R}^n; \Lambda^k\mathbf{R}^n)$ and let $\Sigma \subset \Omega \subset \mathbf{R}^n$ be open and bounded. Since Σ has vanishing capacity, there is a sequence $(\eta_m)_{m \geq 1}$ in $\mathcal{D}(\Omega)$ such that $0 \leq \eta_m \leq 1$, $\eta_m = 1$ on a neighborhood of Σ and $\|\nabla\eta_m\|_{L^n(\mathbf{R}^n)} \rightarrow 0$ as $n \rightarrow \infty$. Moreover, by Poincaré’s inequality, up to a subsequence, $\eta_m \rightarrow 0$ almost everywhere.

Consider now the sequence

$$\psi_m = (1 - \eta_m)\zeta_m\psi,$$

where ζ_m is given by Lemma A.4. By definition, $\psi_m \in \mathcal{D}(\mathbf{R}^n; \Lambda^{k-1}\mathbf{R}^n)$. We claim that $d\psi_m \rightarrow \varphi$ in $L^1(\mathbf{R}^n; \Lambda^k\mathbf{R}^n)$.

In fact,

$$d\psi_m = -\zeta_m d\eta_m \wedge \psi + (1 - \eta_m) d\zeta_m \wedge \psi + (1 - \eta_m) \zeta_m \varphi. \tag{A.1}$$

By Hölder’s inequality,

$$\|-\zeta_m d\eta_m \wedge \psi\|_{L^1(\mathbf{R}^n)} \leq \|\zeta_m\|_{L^\infty(\mathbf{R}^n)} \|d\eta_m\|_{L^n(\mathbf{R}^n)} \|\psi\|_{L^{n/(n-1)}(\mathbf{R}^n)}.$$

Since $\|\nabla\eta_m\|_{L^n(\mathbf{R}^n)} \rightarrow 0$ and $\|\psi\|_{L^{n/(n-1)}(\mathbf{R}^n)} < \infty$, the first term in (A.1) tends to zero. A similar reasoning holds for the second term, and the last term converges to φ as $m \rightarrow \infty$ by Lebesgue’s dominated convergence theorem. \square

Corollary A.7. *The set $\mathcal{D}_{\#}(\mathbf{R}^n; \Lambda^k\mathbf{R}^n)$ is dense in $L^1_{\#}(\mathbf{R}^n; \Lambda^k\mathbf{R}^n)$.*

A.2. The closure of exact n -forms

Theorem A.6 fails when $k = n$, and therefore the proof Theorem A.5 fails in this case, but there is in fact a stronger result.

Theorem A.8. *The set $d(\mathcal{D}(\mathbf{R}^n; \Lambda^{n-1}\mathbf{R}^n))$ is dense in $\mathcal{D}_{\#}(\mathbf{R}^n; \Lambda^n\mathbf{R}^n)$.*

Remark A.9. The density is with respect to the usual topology on the space of test functions [16].

Proof of Theorem A.8. Let $\varphi \in \mathcal{D}_\#(\mathbf{R}^n; \Lambda^n \mathbf{R}^n)$. Therefore $\varphi = f \omega_1 \wedge \dots \wedge \omega_n$, with $f \in \mathcal{D}(\mathbf{R}^n)$ and $\int_{\mathbf{R}^n} f \, dx = 0$. Let $(\rho_\varepsilon)_{\varepsilon>0}$ be a sequence of mollifiers. Define $g_\varepsilon \in \mathcal{D}(\mathbf{R}^n; \mathbf{R}^n)$ by

$$g_\varepsilon(z) = \frac{2}{\|f\|_{L^1(\mathbf{R}^n)}} \int_{\mathbf{R}^n \times \mathbf{R}^n} (x - y) \int_0^1 \rho_\varepsilon(z - tx - (1-t)y) f_+(x) f_-(y) \, dt \, dx \, dy.$$

Next, note

$$\begin{aligned} \operatorname{div} g_\varepsilon(z) &= \frac{2}{\|f\|_{L^1(\mathbf{R}^n)}} \int_{\mathbf{R}^n \times \mathbf{R}^n} \int_0^1 (x - y) \cdot \nabla \rho_\varepsilon(z - tx - (1-t)y) f_+(x) f_-(y) \, dt \, dx \, dy \\ &= \frac{2}{\|f\|_{L^1(\mathbf{R}^n)}} \int_{\mathbf{R}^n \times \mathbf{R}^n} (\rho_\varepsilon(z - x) - \rho_\varepsilon(z - y)) f_+(x) f_-(y) \, dx \, dy \\ &= (\rho_\varepsilon * f)(z). \end{aligned} \tag{A.2}$$

Therefore $\operatorname{div} g_\varepsilon \rightarrow f$ in $\mathcal{D}(\mathbf{R}^n)$ as $\varepsilon \rightarrow 0$. Letting

$$\psi_\varepsilon = \sum_{i=1}^n g_\varepsilon^i (-1)^{i+1} \omega_1 \wedge \dots \wedge \widehat{\omega}_i \wedge \dots \wedge \omega_n,$$

one concludes

$$d\psi_\varepsilon = \operatorname{div} g_\varepsilon \omega_1 \wedge \dots \wedge \omega_n \rightarrow f \omega_1 \wedge \dots \wedge \omega_n = \varphi$$

as $\varepsilon \rightarrow 0$. \square

Remark A.10. The construction (A.2) is inspired from the construction of a non-optimal mass displacement plan in the Monge–Kantorovich mass displacement problem [11].

References

[1] R.A. Adams, Sobolev Spaces, Pure Appl. Math., vol. 65, Academic Press, New York, 1975.
 [2] F. Bethuel, G. Orlandi, D. Smets, Approximations with vorticity bounds for the Ginzburg–Landau functional, *Commun. Contemp. Math.* 6 (5) (2004) 803–832.
 [3] J. Bourgain, H. Brezis, New estimates for the Laplacian, the div–curl, and related Hodge systems, *C. R. Math. Acad. Sci. Paris* 338 (7) (2004) 539–543.
 [4] J. Bourgain, H. Brezis, P. Mironescu, $H^{1/2}$ maps with values into the circle: Minimal connections, lifting, and the Ginzburg–Landau equation, *Publ. Math. Inst. Hautes Études Sci.* 99 (2004) 1–115.
 [5] H. Brezis, *Analyse fonctionnelle*, Collect. Math. Appl. Maîtrise, Masson, Paris, 1983.
 [6] H. Brezis, L. Nirenberg, Degree theory and BMO. I. Compact manifolds without boundaries, *Selecta Math. (N.S.)* 1 (2) (1995) 197–263.
 [7] H. Brezis, J. Van Schaftingen, L^1 estimates on domains, in preparation.
 [8] D.-C. Chang, G. Dafni, E.M. Stein, Hardy spaces, BMO, and boundary value problems for the Laplacian on a smooth domain in \mathbf{R}^n , *Trans. Amer. Math. Soc.* 351 (4) (1999) 1605–1661.

- [9] L.C. Evans, R.F. Gariepy, *Measure theory and fine properties of functions*, Stud. Adv. Math., CRC Press, Boca Raton, FL, 1992.
- [10] H. Federer, *Geometric Measure Theory*, Springer-Verlag, New York, 1969.
- [11] L. Kantorovitch, G. Akilov, *Analyse fonctionnelle. Tome 1. Opérateurs et fonctionnelles linéaires*, Mir, Moscow, 1981.
- [12] C.J. Neugebauer, Lipschitz spaces and exponentially integrable functions, *Indiana Univ. Math. J.* 23 (1973–1974) 103–106.
- [13] H.M. Reimann, Functions of bounded mean oscillation and quasiconformal mappings, *Comment. Math. Helv.* 49 (1974) 260–276.
- [14] T. Runst, W. Sickel, *Sobolev spaces of Fractional Order, Nemytskij Operators, and Nonlinear Partial Differential Equations*, de Gruyter Ser. Nonlinear Anal. Appl., vol. 3, de Gruyter, Berlin, 1996.
- [15] D. Sarason, Functions of vanishing mean oscillation, *Trans. Amer. Math. Soc.* 207 (1975) 391–405.
- [16] L. Schwartz, *Théorie des distributions*, vols. IX–X, Publ. Inst. Math. Univ. de Strasbourg, Hermann, Paris, 1966.
- [17] L. Simon, *Lectures on Geometric Measure Theory*, Proc. Centre Math. Anal. Austral. Nat. Univ., vol. 3, Australian National University, Centre for Mathematical Analysis, Canberra, 1983.
- [18] S.K. Smirnov, Decomposition of solenoidal vector charges into elementary solenoids, and the structure of normal one-dimensional flows, *Algebra i Analiz* 5 (4) (1993) 206–238.
- [19] E.M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Math. Ser., vol. 30, Princeton University Press, Princeton, NJ, 1970.
- [20] E.M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Math. Ser., vol. 43, Princeton University Press, Princeton, NJ, 1993.
- [21] H. Triebel, *Interpolation Theory, Function Spaces, Differential Operators*, North-Holland Math. Library, vol. 18, North-Holland, Amsterdam, 1978.
- [22] J. Van Schaftingen, A simple proof of an inequality of Bourgain, Brezis and Mironescu, *C. R. Math. Acad. Sci. Paris* 338 (1) (2004) 23–26.
- [23] J. Van Schaftingen, Estimates for L^1 -vector fields, *C. R. Math. Acad. Sci. Paris* 339 (3) (2004) 181–186.
- [24] W.P. Ziemer, *Weakly Differentiable Functions: Sobolev Spaces and Functions of Bounded Variation*, Grad. Texts in Math., vol. 120, Springer-Verlag, New York, 1989.