

On the lowest eigenvalue of a pseudo-differential operator

(sharp Gårding inequalities/uncertainty principle/subelliptic estimates/commutators of vector fields)

C. FEFFERMAN* AND D. H. PHONG†

*Department of Mathematics, Princeton University, Princeton, New Jersey 08544; and †Department of Mathematics, Columbia University, New York, New York 10027

Contributed by Charles Fefferman, September 17, 1979

ABSTRACT Positive lower bounds for pseudo-differential operators with nonnegative symbols are derived; the bounds in particular yield subelliptic estimates for operators arising as sums of squares of vector fields.

Let $p(x, \xi)$ be a nonnegative symbol satisfying the estimates

$$|D_x^\alpha D_\xi^\beta p(x, \xi)| \leq C_{\alpha\beta} M^{2-|\beta|}. \quad [1]$$

We shall outline an algorithm to determine the order of magnitude of the lowest eigenvalue of the corresponding pseudo-differential operator $p(x, D)$. This is closely related to earlier work on conditions ensuring the estimate

$$\operatorname{Re} \langle p(x, D)u, u \rangle + C \|u\|^2 \geq 0 \quad u \in L^2(\mathbb{R}^n). \quad [2]$$

The sharpest known sufficient conditions for inequality 2 are the following:

- (i) $p \in S^2(\mathbb{R}^n \times \mathbb{R}^n)$, $p \geq 0$ (see ref. 1)
- (ii) $p \in S^{6/5}(\mathbb{R}^n \times \mathbb{R}^n)$, $p + \operatorname{Tr}^+ p \geq 0$, in which

$\operatorname{Tr}^+ p$ is a nonnegative quantity defined in terms of the Hessian of p [see Hörmander (2) and also Melin (3)].

Our first main result on the eigenvalue problem, motivated by the uncertainty principle of quantum mechanics, is the following:

Let $Q_0 = \{(x, \xi) \in \mathbb{R}^n \times \mathbb{R}^n; |x|, |\xi| \leq 1\}$; say that a canonical transformation $\Phi: (x, \xi) \rightarrow (y, \eta)$ mapping Q_0 into $\mathbb{R}^n \times \mathbb{R}^n$ is a testing map if $y - y_0$ and $(\eta - \eta_0)/M$ are C^α functions of (x, ξ) with norms bounded by a fixed constant. Here (y_0, η_0) denotes $\Phi(0, 0)$, and α is an integer that depends on ε below.

THEOREM 1. If $p(x, \xi) \geq 0$ satisfies inequality 1, and $K \geq C_\varepsilon M^\varepsilon$ is a constant such that

$$\|p \circ \Phi\|_{C^0(Q_0)} \geq K \text{ for any testing map } \Phi,$$

then

$$\operatorname{Re} \langle p(x, D)u, u \rangle \geq c_\varepsilon K \|u\|^2 \quad u \in L^2(\mathbb{R}^n). \quad [3]$$

From Theorem 1, one can easily read off the following special case of the theorem of Hörmander (4) on commutators of vector fields:

COROLLARY. Let X_1, \dots, X_m be vector fields on \mathbb{R}^n whose Lie brackets up to order k generate the Lie algebra at each point. Then

$$\sum_{j=1}^m \|X_j u\|^2 + C_\varepsilon \|u\|^2 \geq c_\varepsilon \|u\|^{2-\varepsilon+1/(k+1)} \quad \varepsilon > 0 \quad [4]$$

for $u \in C^\infty$ supported in the unit ball in \mathbb{R}^n .

In fact, inequality 4 holds for $\varepsilon = 0$, as was proved by Rothschild and Stein (5) (together with estimates in norms other than L^2); we shall also derive that result from a refinement of Theorem 1 to be given below.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U. S. C. §1734 solely to indicate this fact.

Proof of the Corollary. First observe that

$$\|f\|_{C^k(Q_0)} \leq C_\varepsilon M^\varepsilon \|f\|_{C^0(Q_0)} + C'_\varepsilon M^{-2} \|f\|_{C^0(Q_0)} \quad [5]$$

if $\beta > [3k/\varepsilon] + 1$, as can be seen by comparing f with its Taylor expansion of order β in a ball of radius $M^{-\varepsilon/k}$. Next let p_1, \dots, p_m be the symbols of X_1, \dots, X_m localized to a conic neighborhood U in the region $|\xi| \approx M$, and assume that

$$|\{p_{i_1}, \{p_{i_2}, \dots, \{p_{i_k}, p_{i_{k+1}}, \dots\}\}\}| \approx M \quad \text{throughout } U. \quad [6]$$

If Φ is a testing map then expression 6 will still hold with p_{i_j} replaced by $p_{i_j} \circ \Phi$. The commutator is of course a polynomial homogeneous of degree $k + 1$ in derivatives (of order up to k) of the $p_{i_j} \circ \Phi$, and therefore expression 6 yields

$$\|p_{i_j} \circ \Phi\|_{C^k(Q_0)} \geq cM^{1/k+1}$$

for some i_j , which in turn implies

$$\max_j \|p_{i_j} \circ \Phi\|_{C^0(Q_0)} \geq c_\varepsilon M^{-\varepsilon+1/(k+1)}$$

in view of inequality 5. The corollary is now an immediate consequence of Theorem 1.

The proof of Theorem 1 proceeds by induction on the number of variables; a key tool in the induction is the following theorem, which also provides an algorithm for computing the size of the lowest eigenvalue:

THEOREM 2. Suppose $p(x, \xi) \geq 0$ satisfying inequality 1 in $\{|\xi_n| < M\}$ is of the form

$$p(x, \xi) = \xi_n^2 + a(x, \xi'), \text{ in which } \xi' = (\xi_1, \dots, \xi_{n-1}). \quad [7]$$

We define its derived symbol to be

$$p^*(x', x_n, \xi') = \sum_{j=0}^N a(x', x_n + jK^{-1/2}/N, \xi'),$$

in which $N = [3/\varepsilon] + 1$. Then inequality 3 is equivalent to

$$\operatorname{Re} \langle p^*(x', x_n, D')v, v \rangle_{L^2(\mathbb{R}^{n-1})} \geq cK \|v\|_{L^2(\mathbb{R}^{n-1})}^2 \quad [8]$$

for each x_n in \mathbb{R} and v in $L^2(\mathbb{R}^{n-1})$.

We now present the promised algorithm, which will either determine the lowest eigenvalue λ of $p(x, D)$ up to a bounded factor or else show that $\lambda \leq C_\varepsilon M^\varepsilon$. Assume we already know how to carry out our algorithm in $(n - 1)$ dimensions ($n \geq 1$); given a symbol $p \geq 0$ on $\mathbb{R}^n \times \mathbb{R}^n$ satisfying inequality 1, and a constant $K \geq C_\varepsilon M^\varepsilon$, our task is to decide whether the estimate 3 holds. Perform a Calderón-Zygmund decomposition of the basic $1 \times M$ blocks in $\mathbb{R}^n \times \mathbb{R}^n$, stopping at Q_s if one of the following occurs:

$$(i) \quad (\min_{Q_s} p)(\operatorname{diam}_x Q_s)^{-2}(\operatorname{diam}_\xi Q_s)^{-2} \geq A,$$

$$(ii) \quad \max_{|\alpha|+|\beta|=2} \|D_x^\alpha D_\xi^\beta p\|_{L^\infty(Q_s)} \times (\operatorname{diam}_x Q_s)^{|\alpha|-2}(\operatorname{diam}_\xi Q_s)^{|\beta|-2} \geq A,$$

$$(iii) \quad (\text{diam}_x Q_s)(\text{diam}_\xi Q_s) \leq K^{1/2}/A.$$

Here A is a large constant, and Q_s^* is the dilate of Q_s by a large constant factor. In view of the $S_{\phi, \varphi}^{M, m}$ calculus (see ref. 6), inequality 3 holds for $p(x, \xi)$ if and only if localized estimates hold for $p|_{Q_s}$. Thus inequality 3 is evidently false if there is any Q_s satisfying iii. Otherwise, because the localized estimate is obviously true for Q_s satisfying i, the only delicate case is ii. However, a suitable canonical transformation carries the symbol $p|_{Q_s}$ to a symbol of the form 7, so that *Theorem 2* reduces the problem to an eigenvalue computation in fewer variables.

The estimate 4 with $\varepsilon = 0$ can be obtained from our algorithm, which in fact shows that if $p = \sum p_j^2$ and

$$|\{p_{i_1}, \{p_{i_2}, \dots, \{p_{i_l}, p_{i_{l+1}}\} \dots}\}| \geq K^{(l+1)/2} \text{ for some } l, \quad [9]$$

then inequality 3 holds, the reason being essentially that the derived symbol of p arising from a cube Q_s of type ii is again a sum of squares satisfying hypotheses analogous to inequality 9.

Theorem 2 in turn can be deduced from the following result on the spectral decomposition of pseudo-differential operators, which may be of intrinsic interest:

THEOREM 3. Given $p(x, \xi) \geq 0$ and a constant K , let

$$p_K(u) = \langle \min\{K, p^w(x, D)\}u, u \rangle,$$

in which $p^w(x, D)$ is defined by the Weyl calculus as in Hörmander (2), and the minimum is taken in the sense of spectral theory. Then if p, q are nonnegative symbols satisfying inequality 1, we have

$$(p + q)_K(u) \leq C_\varepsilon [p_K(u) + q_K(u) + M^\varepsilon \|u\|^2].$$

The proofs of the results announced here will appear in a forthcoming article.

It would be interesting to know whether the lower bound for the least eigenvalue of $p(x, D)$ given by *Theorem 1* is sharp.

This research was carried out while C.F. was supported by the National Science Foundation Alan T. Waterman Award and D.H.P. was partially supported by the National Science Foundation.

1. Fefferman, C. & Phong, D. H. (1978) *Proc. Natl. Acad. Sci. USA* **75**, 4673–4674.
2. Hörmander, L. (1979) *Commun. Pure Appl. Math.* **32**, 359–443.
3. Melin, A. (1971) *Ark. Mat.* **9** 117–140.
4. Hörmander, L. (1967) *Acta Math.* **119**, 147–171.
5. Rothschild, L. & Stein, E. M. (1977) *Acta Math.* **137**, 247–320.
6. Beals, R. & Fefferman, C. (1974) *Commun. Pure Appl. Math.* **27**, 1–24.